CORRECTED VERSION*

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶:
C12N 15/12, 15/85, 5/10, C07K 14/705, 16/28, C12Q 1/68, A61K 38/17, 48/00

(11) International Publication Number:

WO 98/31799

(43) International Publication Date:

23 July 1998 (23.07.98)

(21) International Application Number:

PCT/US98/00959

A3

(22) International Filing Date:

21 January 1998 (21.01.98)

(30) Priority Data:

60/034,204 60/034,205 21 January 1997 (21.01.97) US

21 January 1997 (21.01.97) US

(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). GENTZ, Reiner, L. [DE/US]; 13404 Fairland Park Drive, Silver Spring, MD 20904 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US).
- (74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(88) Date of publication of the international search report: 5 November 1998 (05.11.98)

- (54) Title: POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS
- (57) Abstract

Receptor polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing receptor polypeptides and polynucleotides in the design of protocols for the treatment of diseases and diagnostic assays for such conditions.

FOR THE PURPOSES OF INFORMATION ONLY

en de junta segende el entre o de la completa esta esta esta entre esta en entre esta en especial esta entre e

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ.	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	ΙE	ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	iceland	MW	Malawi	us	United States of Americ
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KĠ	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

WO 98/31799 PCT/US98/00959

POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

FIELD OF INVENTION

5

10

15

20

25

30

This invention relates to newly identified polynucleotides and the polypeptides encoded by them, the use of such polynucleotides and polypeptides, and their production. More particularly, the polynucleotides and polypeptides of the present invention relate to specific receptor families described in the specification and known in the art. The invention also relates to inhibiting or activating the action of such polynucleotides and polypeptides.

BACKGROUND OF THE INVENTION

Receptor proteins are found on the membrane of the cells and are generally involved in signal transduction. There are many types of receptor proteins, and for convenience, these proteins are grouped in families based on similarity in structure and function.

For example, the TM4SF superfamily of cell surface proteins, also known as the tetraspan receptor superfamily, is comprised of at least seventeen individual gene products (these include CD9, CD20, CD37, CD53, CD63, CD81, CD82, A15, CO-029, Sm23, RDS, Uro B, Uro A, SAS, Rom-1, PETA3, and YKK8). The TM4SF superfamily is the second largest group in the CD antigen superfamily. Each member of the TM4SF superfamily can be characterized by several putative physical features including four highly conserved transmembrane domains, two divergent extracellular loops, and two short and highly divergent cytoplasmic tails. Expression patterns for members of the TM4SF superfamily tend to be rather broad and can vary widely between members. The functional roles of TM4SF superfamily members are primarily associated with signal transduction events and pathways, but also include cell adhesion in platelets and other lymphocytic and non-lymphocytic cell lines, as well as cell motility, proliferation, and metastasis. In addition, recent evidence suggests that a subset of the members of the TM4SF superfamily may function as potassium channel molecules.

One member of the TM4SF family, CD20, is a four membrane spanning domain cell surface phosphoprotein expressed exclusively on B lymphocytes.

35 Although the precise functional role of CD20 has yet to be determined, it is thought to function primarily as a receptor during B-cell activation. Furthermore, a large number of experimental observations suggest several additional speculative roles for the CD20 molecule. For example, CD20-specific immunoprecipitation of biochemically cross-linked plasma membrane proteins suggests that CD20 assumes a multimeric structural

WO 98/31799 2 PCT/US98/00959

conformation characteristic of other previously described membrane channel proteins. Further experimentation has revealed that expression of exogenous CD20 on the cell surface specifically increases Ca²⁺ conductance across the plasma membrane. Together, these results suggest that CD20 complexes may function as B-cell specific Ca²⁺ ion channels. In addition, monoclonal antibodies raised against CD20 have been used to stimulate resting B-cells to transition out of the G0/G1 segment of the cell cycle. It has also been demonstrated that CD20 is associated with both serine and tyrosine kinases and, more specifically, that CD20 is associated, although not directly, with the Src family of tyrosine kinases including p56/53lyn, p56lck, and p59fyn.

A second example of a receptor subfamily, called sialoadhesin molecules, belongs to the Ig superfamily of receptor-like molecules. The more than 100 members of the Ig superfamily are generally considered to engage in specific cell-cell interactions through which intercellular communication may occur. In addition to classical protein-protein interactions, intercellular communication may also be mediated through protein-carbohydrate interactions. In fact, all members of the sialoadhesin family of the Ig superfamily are capable of mediating protein-sialic acid binding interactions. To date, only a small number of proteins have been assigned to the sialoadhesin family including sialoadhesin, CD33, CD22, the myelin-associated glycoprotein (MAG), and the Schwann cell myelin protein (SMP). Each of these proteins is expressed in a restricted subset of cell types. For example, CD22 and CD33 are expressed exclusively by B-lymphocytes and cells of the myelomonocytic lineage, respectively.

Similarly, galectins are a family of the lectin superfamily of carbohydrate-binding proteins which have a high affinity for b-galactoside sugars. Although a large number of glycoproteins containing b-galactoside sugars are produced by the cell, only a few will bind to known galectins *in vitro*. Such apparent binding specificity suggests a highly specific functional role for the galectins. Galectin 1 (conventionally termed LGALS1 for lectin, galactoside-binding, soluble -1) is thought to specifically bind laminin, a highly polylactosaminated cellular glycoprotein, as well as the highly polylactosaminated lysosome-associated membrane proteins (LAMPs). Galectin 1 has also been shown to bind specifically to a lactosamine-containing glycolipid found on olfactory neurons and to integrin a_7b_1 on skeletal muscle cells. Galectin 3 has also been observed to bind specifically to laminin, immunoglobulin E and its receptor, and bacterial lipopolysaccharides.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin 1 appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1 mediated disruption of laminin-integrin a_7b_1 interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined

5

10

15

20

25

30

WO 98/31799 3 PCT/US98/00959

presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bc12, respectively.

A relatively new family of cell-surface proteins has been identified and termed the Ly6 superfamily. The members of this family include murine and human SCA-2, rat Ly-6 (also termed ThB), human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)], and E48 antigen. The determination of an initial functional role for SCA-2 may lie in an analysis of its expression profile with regard to the complex process of hematopoiesis. SCA-2 is highly expressed in early thymic precusor cells. In turn, progeny of the intrathymic precusor population continue to express SCA-2, but only until the point of transition occurs from blast cell to small cell. Further experimental evidence demonstrates that mature thymocytes and peripheral T-cells do not express detectable levels of SCA-2, whereas mature, peripheral B-cells do continue to express SCA-2. As a result, it seems very likely that SCA-2 plays an important role in thymocyte maturation and differentiation. A plausible explanation for this functional hypothesis is that SCA-2 may act as a receptor for a unknown cytokine which regulates thymocyte maturation and differentiation.

In addition, CD59 is a recently identified integral membrane protein which appears to be involved in the regulation of complement. Recent studies show that the CD59 antigen may prevent damage from complement C5b-9 and protect astrocytes during inflammatory and infectious disorders of the nervous system. Expression of recombinant human CD59 on porcine donor organs have been shown to prevent complement-mediated lysis and activation of endothelial cells that leads to hyperacute rejection. Recently, researchers at Alexion Pharmaceuticals (New Haven, CT) reported on the production of transgenic pigs which expressed human CD59. In these animals, xenogeneic organs were resistant to hyperacute rejection. (Fodor, et al., "Expression of a functional human complement inhibitor in a transgenic pig as a model for the prevention of xenogeneic hyperacute organ rejection," Proc. Natl. Acad. Sci., 91:1153-11157 (1994).) The same company also reported that expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria (PNH) B-

WO 98/31799 4 PCT/US98/00959

and the state of t

cells confers resistance to human complement. (Rother et al., "Expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria B-cells confers resistance to human complement," Blood, 84:2604-2611 (1994).) PNH is an acquired hematopoietic disorder characterized by complement-mediated hemolytic anemia, pancytopenia, and venous thrombosis. It is thought that retroviral gene therapy with this molecule could provide a treatment for PNH patients.

A final Ly6 superfamily member, the E48 antigen, is involved in intercellular adhesion between keratinocyte cells of the squamous epithelium. Such keratinocytes are attached to adjoining cells by large numbers of desmosomes, which are thought to play a role in the transition of transformed keratinocytes to metastatic tumor cells. Treatment with a monoclonal antibody raised against the E48 antigen has been successful in the eradication of residual, postoperative squamous cell carcinoma cells of the upper aerodigestive tract in several *in vivo* models and, to some degree, in humans. (van Dongen, et al., "Progress in radioimmunotherapy of head and neck cancer," Oncol. Rep. 1:259-264 (1994).) The gene encoding the E48 antigen has been mapped to the q24-qter region of human chromosome 8. Interestingly, a number of human diseases have been mapped to this region of chromosome 8 including Langer-Giedion syndrome, brachio-otorhinolaryngeal syndrome, trichorhinolaryngeal syndrome, and epidermolysis bullosa simplex.

A further example of a receptor family includes the prohibitin receptors. The prohibitin gene product is expressed in a wide variety of tissues and has been implicated as a component of a number of anti-proliferative mechanisms. The prohibitin gene encodes a 30 kD postsynthetically modified polypeptide located primarily in the mitochondria, but also may be associated with the IgM receptor on the B-cell plasma membrane. The protein functionally inhibits DNA synthesis and entry into S phase of the cell cycle by an unknown mechanism. Interestingly, although the prohibitin gene product is hypothesized to be involved in the maintenance of senescence and the prevention of cancer, one study found that, although somatic mutations in the prohibitin gene were present in a small number of breast cancers, no mutations were identified in any other breast, ovary, liver, and lung cancers examined. (Sato et al., Genomics 17:762-764 (1993).) However, the prohibitin gene has been mapped to human chromosome 17q12-21, the same region thought to contain the gene involved in sporadic breast cancer. Furthermore, DNA sequence analysis of the prohibitin gene identified somatic mutation in 4 of 23 cases of sporadic breast cancer examined. Thus, prohibitin family members may be involved in the development of cancer.

Moreover, the EGFR family of plasma membrane proteins are an integral component of normal cellular proliferation and in the pathogenesis of the cancerous state. The family is relatively small and includes the EGFR, c-erbB-2, c-erbB-3, and others. Various cancers are correlated with aberrant expression of one or more of these

5

10

15

20

25

30

10

15

20

25

30

35

genes. A number of ligands have been identified which bind to the EGFR-like receptors listed above including TGF-a, heparin-binding EGF, amphiregulin, criptoregulin, heregulin, and others. A large fraction of adenocarcinomas examined to date, especially those of the breast, colon, and pancreas, are typified by the amplification or overexpression of the c-erbB-2 gene. EGF, or an analogous ligand, initiates the cellular growth factor response by binding to the EGFR, or EGFR-related, receptor. Following the binding event, the receptor molecule dimerizes activating its intracellular tyrosine kinase domain. This event results in the phosphorylation of specific tyrosine residues near the carboxy terminus of the receptor. The diversity of signals able to be transduced through the relatively small number of EGFR-related receptor molecules is amplified considerably by the recent finding that EGFR-like receptor molecules can function when dimerized with other EGFR family members forming heterodimers.

Members of the EGFR-related family of integral membrane proteins have been implicated in the pathogenesis of a number of human disease-states. For example, a mutation in the EGFR itself appears to play an important role in the development of glioblastomas. (Sang et al., J. Neurosurg 82:841-846 (1995).) The EGFR gene is amplified or overexpressed in the majority of primary human glioblastomas. Although not conferring a distinct advantage on cell growth, an increase in EGFR expression was found to confer an increase in the ability of glioma cells to maintain anchorage-independent growth in soft agar especially in response to EGF and retinoic acid. Anchorage-independent growth *in vitro* correlates highly with tumorigenicity *in vivo*, therefore, it is likely that cells which express abnormally high levels of EGFR in human glioblastoma cells may be involved in the high potential for these cells to cause tumors *in vivo*.

Moreover, overexpression or amplification of c-erbB-2 has been reported to be involved in a high number adenocarcinomas, particularly of the breast, colon, and pancreas, and in a small proportion of ovarian carcinomas.

Thus, there is a clear need for identifying and exploiting novel members of the receptor families, such as those described above. Although structurally related, these receptors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such receptors may prove useful as therapeutics in an antitumor, diagnostic, or other capacity. Furthermore, receptors described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

WO 98/31799 6 PCT/US98/00959

SUMMARY OF THE INVENTION

5

10

15

20

25

30

35

in a construction of the control of

In one aspect, the invention relates to receptor polypeptides and polynucleotides, as well as the methods for their production. Another aspect of the invention relates to methods for using such receptor polypeptides and polynucleotides. Such uses include the treatment of the specified diseases, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with receptor imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate receptor activity or levels.

DESCRIPTION OF THE INVENTION Definitions

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Receptor" refers, among others, to a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:Y, or an allelic variant thereof.

"Receptor Activity" or "Biological Activity of the Receptor" refers to the metabolic or physiologic function of said receptor including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said receptor.

"Receptor gene" refers to a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO:X or allelic variants thereof and/or their complements.

"SEQ ID NO:X" comprises all or a substantial portion of the polynucleotide encoding each receptor of the invention. The value X for the nucleotide sequence is an integer specified in Table 1. This nucleotide sequence was translated into the receptor polypeptide identified in Table 1 as "SEQ ID NO:Y," where the value of Y for each receptor polypeptide is an integer defined in Table 1.

The invention further provides a composition of matter comprising a nucleic acid molecule which comprises a human cDNA clone identified by a cDNA Clone ID (Identifier) in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection ("ATCC") and given the ATCC Deposit Number shown in Table 1 for that cDNA clone. The ATCC is located at American Type Culture Collection (ATCC), 12301 Park Lawn Drive. Rockville, Maryland 20852, USA. The deposit has been made under the terms of the Budapest Treaty on the international recognition of the deposit of micro-organisms for purposes of patent procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposit is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that

WO 98/31799 7 PCT/US98/00959

required under 35 U.S.C. §112. The nucleotide sequence of the polynucleotides contained in the deposited material, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

5

10

15

20

25

30

35

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can

kinder og er tokkallista komzettil blevet hved littligt i helle sig etter i skrivet i skrivet i skrivet i skri

WO 98/31799 8 PCT/US98/00959

occur anywhere in a polypeptide, including the peptide backbone, the amino acid sidechains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphotidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., "Analysis for protein modifications and nonprotein cofactors", Meth Enzymol (1990) 182:626-646 and Rattan et al., "Protein Synthesis: Posttranslational Modifications and Aging", Ann NY Acad Sci (1992) 663:48-62.)

"Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions; identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to

5

10

15

20

25

30

occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" per se has an art-recognized meaning and can be calculated using published techniques. (See, e.g.: COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; 10 SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991.) While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans. (Carillo, H., and Lipton, D., SIAM J 15 Applied Math (1988) 48:1073.) Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred 20 computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCS program package (Devereux, J., et al., Nucleic Acids Research (1984) 12(1):387), BLASTP, BLASTN, FASTA (Atschul, S.F. et al., J Molec Biol (1990) 215:403.)

25

30

35

As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO:X is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: X. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5 ' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

WO 98/31799 10 PCT/US98/00959

Same of the same of the same

Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% "identity" to a reference amino acid sequence of SEQ ID NO:Y is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO:Y. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

15

20

25

10

5

Polypeptides of the Invention

In one aspect, the present invention relates to receptor polypeptides (or receptor proteins). The receptor polypeptides include the polypeptide of SEQ ID NO:Y; as well as polypeptides comprising the amino acid sequence of SEQ ID NO:Y; and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% identity to SEQ ID NO:Y are highly preferred. Also included within receptor polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% are highly preferred. Preferably receptor polypeptides exhibit at least one biological activity of the receptor.

30

The receptor polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, prosequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

35

Fragments of the receptor polypeptides are also included in the invention. A "fragment" is a polypeptide having an amino acid sequence that entirely is the same as part, but not all, of the amino acid sequence of the aforementioned receptor polypeptides. As with receptor polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most

10

15

20

25

30

35

San San Francis

preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of receptor polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of receptor polypeptides, except for deletion of a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus.

Also preferred are fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. The "domains" of each receptor polypeptide are illustrated in the Figures. The Figures compare SEQ ID NO:Y to the closest know homologue. Identical amino acids shared between the two polypeptides are shaded, while conservative amino acid changes are boxed. By examining the regions or amino acids shaded and/or boxed, the skilled artisan can readily identify conserved domains between the two polypeptides. The amino acids sequences of SEQ ID NO:Y falling within these conserved domains are "fragments" and are specifically contemplated by the present invention. Especially preferred is the extracellular domains of a receptor of the invention. Soluble extracellular domains have antagonist activity mediated by competition with a receptor ligand.

Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate receptor activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

Preferably, all of these polypeptide fragments retain a biological activity of the receptor, including antigenic activity. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions -- i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

WO 98/31799 12 PCT/US98/00959

The receptor polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

Polynucleotides of the Invention

5

10

15

20

25

30

35

Another aspect of the invention relates to receptor polynucleotides. Receptor polynucleotides include isolated polynucleotides which encode the receptor polypeptides and fragments, and polynucleotides closely related thereto. More specifically, a receptor polynucleotide of the invention includes a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:X encoding a receptor polypeptide of SEQ ID NO:Y, and polynucleotide having the particular sequence of SEQ ID NO:X.

Receptor polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y, and a polynucleotide comprising a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under receptor polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO:X, or contained in the cDNA insert in the plasmid deposited with ATCC, to hybridize under conditions useable for amplification or for use as a probe or marker. Moreover, the receptor polynucleotide includes a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC, and a nucleotide sequence comprising at least 15 contiguous nucleotides of such cDNA insert. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. The invention also provides polynucleotides which are complementary to all the above receptor polynucleotides.

The receptors of the invention are structurally related to other proteins of specified receptor families, as shown by the results in the Figures. The cDNA sequence of SEQ ID NO:X encodes a polypeptide as described in Table 1 as SEQ ID NO:Y. Because the receptor polypeptides contain domains similar in structure to other receptor family members, the receptors of the present invention are expected to have,

inter alia, similar biological functions/properties to their homologous polypeptides and polynucleotides, and their utility is obvious to anyone skilled in the art.

Table 1

5

10

15

Clone ID	SEQ	SEQ	ATCC	ATCC	Receptor	Homology
Name	ID NO:X	ID NO:Y	Deposit	Deposit	Family	
HMACR70	NU:X	NU: Y	No. 209054	Date 05/16/97	la la	Sialoadhesin
HMACK/0	1	18	209034 #####	01/21/98	Ig	OB-1
HTEDK48	1	10	209054	05/16/97	TM4SF	MRC-OX44
HIEDR46			209034	03/10/9/	1 1421.	PETA-3
1-1849 bp	2					
160-900 bp	3	19				
HTPED39			209054	05/16/97	TM4SF	NAG-2
HPWAE25	4	20	#####	1/21/98		TALLA-1
HTPEF86	5	21	209053	05/16/97	TM4SF	CD20
						B1 Antigen
HSBBF02	6	22	209054	05/16/97	TM4SF	TALLA-1
HLTAH80	7	23	97242	08/02/95	TM4SF	TALLA-1
			209054	05/16/97		
HTPBA27	8	24	97242	08/02/95	TM4SF	NAG-2
1			209054	05/16/97	l	
HAIDQ59			209054	05/16/97	TM4SF	CD9
				L		Antigen
5' Sequence	9	25				
3' Sequence	10					
HHFEK40	11	26	209054	05/16/97	TM4SF	PETA-3
HGBGV89	12	27	209125	06/09/97	TM4SF	L6H
			209054	05/16/97		
HUVBB80	13	28	209054	05/16/97	TM4SF	L6
HJACE54	14	29	209053	05/16/97	Lectin	Galectin-3
				,		Galectin-5
				<u> </u>		Galectin-8
HROAD63	15	30	209053	05/16/97	Ly6	E48 splice
						variant
HMWGS46	16	31	209053	05/16/97	Prohibitin	BAP-37
HNFGW06	17	32	209053	05/16/97	EGFR	EGFR

The novel full-length cDNA clone designated HMACR70 may be a member of the sialoadhesin family of the Ig superfamily of receptor-like molecules and a CD33 homologue. HMACR70 contains a 1497 nucleotide cDNA insert encoding a 315 amino acid ORF and was cloned from a GM-CSF-treated human macrophage cDNA library. The only additional cDNA libraries in the HGS database which include this clone are human eosinophils and possibly human gall bladder. A BLAST analysis of the amino acid sequence of HMACR70 demonstrates that this clone exhibits approximately 50% identity and 69% similarity over a 300 amino acids stretch of a gene termed human

WO 98/31799 14 PCT/US98/00959

differentiation antigen, and 38% identity and 62% similarity of the human myelin-associated glycoprotein precursor CD33 gene.

A more recent BLAST analysis confirms HMACR70's designation as a sialoadhesin family member. HMACR70 is homologous to two recently identified sialoadhesin family members, human OB binding protein (OB) 1 and 2. (See, Genbank Accession No. U71382; see Figure 1.) It is thought that OB-1 and OB-2 may bind leptin. Thus, HMACR70, as a sialoadhesin family member, may act to attenuate or even amplify intercellular routes of communication, including binding to leptin or modulating the activity of immune cells, such as macrophages. Clearly, any diseases affected by these processes could be treated by the polypeptide or fragment of HMACR70.

The full-length nucleotide sequences of ten novel human cDNA clones which potentially belong to the TM4SF superfamily are disclosed in the table above and will be addressed sequentially.

The cDNA clone HTEDK48 contains a 1849 nucleotide cDNA insert encoding a 245 amino acid ORF that was cloned from a human testes cDNA library. The coding sequence of HTEDK48 (SEQ ID NO: 3) may be fused to other human proteins, such as 3-hydroxyacyl-CoA dehydrogenase. BLAST analysis of the amino acid sequence of HTEDK48 demonstrates that this clone exhibits approximately 30% identity and 51% similarity over a 245 amino acid stretch of the CD82 molecule. Recent studies have shown that CD82 can associate with CD4 or CD8 and deliver costimulatory signals for the TCR/CD3 pathway. CD82 has also been found to be involved in syncytium formation in HTLV-I-infected T-cells. And finally, in a recently published study in which the expression of the CD82 gene by tumors of the lung was examined retrospectively, it was reported that CD82 may be linked to the suppression of tumor metastasis of prostate cancer. The study also reported that decreased CD82 expression may be involved in malignant progression of such cancers. Thus, HTEDK48 may also be involved in the development of cancer.

A more recent BLAST analysis shows that HTEDK48 is homologous the rat leukocyte antigen, MRC OX-44, and the platelet endothelial tetraspan antigen -3 (PETA-3). (See Figure 2X.) MRC OX-44, a member of a new family of cell surface proteins, appears to be involved in growth regulation. (See, Bellacosa, A., et al., "The Rat Leukocyte antigen MRC OX-44 is a Member of a New Family of Cell Surface Proteins which Appear to be Involved in Growth Regulation," Mol. Cell. Bio. 11: 2864-2872 (1991).) Similarly, PETA-3 has been located to platelet endothelial cells, and an anti-PETA-3 antigen monoclonal antibody can stimulate platelet aggregation and mediator release. (See, Fitter, S., "Molecular Cloning of cDNA Encoding a Novel Platelet-Endothelial Cell Tetra-Span Antigen, PETA-3," Blood, 86(4):1348-1355 (1995).) Thus, HTEDK48 may function similar to MRC OX-44 or PETA-3 to affect

医乳色 化基本 人名西西西 医多种的 医动脉动脉

5

10

15

20

25

30

10

15

20

25

30

35

growth of blood cells. Administering polypeptides or fragments of HTEDK48 may be an effective treatment of blood disorders.

The cDNA clone **HPWAE25** contains a 1288 nucleotide cDNA insert encoding a 273 amino acid ORF that was cloned from a human pancreas tumor cDNA library, while clone **HTPED39** represents a truncated cDNA sequence. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including keratinocytes, ulcerative colitis, striatum depression, lymph node breast cancer, ovarian cancer, stage B2 prostate cancer, kidney medulla, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human cell lines including U937, MM96, WM115, and MDAMB231. A BLAST analysis of the amino acid sequence of HTPED39 demonstrates that this clone exhibits approximately 35% identity and 50% similarity over the entire length of the CD37 molecule. The CD37 antigen is expressed on B cells and on a subpopulation of T cells, but not on pre-B or plasma cells. It has been reported that CD37 expression is downregulated in conjunction with B-cell activation, suggesting that CD37 may be involved in the processes which dictate the activation state of the B-cell.

Moreover, HPWAE25 is also homologous to recently identified TM4SF members, NAG-2 and TALLA-1. (See Figure 3.) NAG-2 is thought to complex with integrins and other TM4SF proteins, while TALLA-1 is a highly specific marker of T-cell acute lymphoblastic leukemia and neuroblastoma. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that Complexs with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997); Takagi, S., "Identification of a Higly Specific Surface Marker of T-cell Acute Lymphoblastic Leukemia and Neuroblastoma as a New Member of the Transmembrane 4 Superfamily," Int. J. Cancer 61(5):706-715 (1995).) Thus, HPWAE25 may be involved the development of cancer, particularly leukemia, lymphoma, and neuroblastoma. HPWAE25 may be used as an effective treatment of these cancers, as well as a diagnostic marker.

A subfamily of TM4SF receptors include CD20 proteins. A CD20-like cDNA clone was obtained from a human pancreas tumor cDNA library and contains a 1236 nucleotide insert which encodes a 250 amino acid ORF. A BLAST analysis of the deduced amino acid sequence of HTPEF86 exhibits approximately 41% identity and 61% similarity to the CD20 gene, also known as B1 antigen. (See Figure 4.) Expression of this gene is detected in only two additional HGS human cDNA libraries; amygdala depression and 9 week early stage human. Although the precise functional role of CD20 has yet to be determined, it is clear that CD20 plays a key role in the regulation of B-cell activation. Based primarily on sequence identity, the novel CD20-like molecule presented herein may also be involved in cell cycle activation. Potential therapeutic and/or diagnostic applications for HTPEF86 may include such clinical

WO 98/31799 16 PCT/US98/00959

presentations as juvenile rheumatoid arthritis, Graves' Disease, and a number of B-cell lymphomas or other lymphoid tumors.

The clone **HSBBF02** contains a 1115 nucleotide cDNA insert encoding a 245 amino acid ORF and was cloned from an HSC 172 cell line cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including brain amygdala depression, endothelial cells, fetal liver and heart, osteoblasts, testes, and others. A BLAST analysis of the amino acid sequence of HSBBF02 demonstrates that this clone exhibits approximately 64% identity and 80% similarity with the A15 molecule over a 131 amino acid stretch (A15 is composed of 244 amino acids). A more recent BLAST search shows that HSBBF02 is similar to the TALLA-1 protein and may in fact be a closely related family member. (See Figure 5.)

In addition, a second cDNA clone, designated **HLTAH80**, exhibits sequence similarity to the A15 molecule and TALLA-1. (See Figure 6.) This clone contains a 1662 nucleotide cDNA insert encoding a 253 amino acid ORF and was cloned from a human T-cell lymphoma cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including B-cell lymphoma, corpus collosum, endometrial tumor, osteosarcoma, testes, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human tissues including spleen, lymph node, thymus, PBLs, heart, and a particularly strong signal in skeletal muscle and pancreas. A BLAST analysis of the amino acid sequence of HLTAH80 demonstrates that this clone exhibits approximately 35% identity and 55% similarity over the entire length of the A15 molecule.

Since expression of A15 drops to undetectable levels when comparing immature T-cells to peripheral blood lymphocytes, it is thought that A15 may play a role in the development of T-cells. Furthermore, the MXS1(CCG-B7) gene which codes for A15 contains a number of triplet nucleotide repeats which have been associated with neuropsychiatric diseases such as Huntington's chorea, fragile X syndrome, and myotonic dystrophy. In addition, A15 appears to be expressed exclusively on T-cell acute lymphoblastic leukemia cell lines, including several derived from adult T-cell leukemia and those established by immortalization with human T-cell leukemia virus type 1 or Herpesvirus saimiri. Thus, clones HLTAH80 and/or HSBBF02 may also be involved in diseases caused by the expansion of repeats or chromosomal instability.

The cDNA clone HTPBA27 contains a 1345 nucleotide cDNA insert encoding a 238 amino acid ORF and was cloned from a human tumor pancreas cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including cerebellum, breast lymph node, osteosarcoma, adult testes, RS4;11 bone marrow cell line, microvascular endothelial cells, and others. A BLAST analysis of the amino acid sequence of HTPBA27 demonstrates that this

5

10

15

20

25

30

WO 98/31799 17 PCT/US98/00959

5

10

15

20

25

30

clone exhibits approximately 40% identity and 64% similarity with a glycoprotein termed CD53 over its entire length. CD53 is thought to be involved in thymopoiesis, since rat CD53 can be detected on immature CD4-8-thymocytes and the functionally mature single-positive subset, but cannot be detected on the intermediate CD4+8+ The CD53 molecule has also been implicated as a thymocytic subset of cells. component of signal transduction pathways in B cells, monocytes and granulocytes, rat macrophages, NK, and T cells. Moreover, as illustrated in Figure 7, HTPBA27 was recently confirmed as a TM4SF receptor. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997).) Calling the HTPBA27 polypeptide NAG-2, this group confirmed HTPBA27's status as a TM4SF receptor by showing that NAG-2 complexes with integrin and other TM4SF receptors. Thus, diseases caused by the failure of HTPBA27 to complex with integrin and other TM4SF receptors can be treated by administering HTPBA27. HTPBA27 can also be used to diagnose these diseases.

The cDNA clone **HAIDQ59** contains cDNA insert encoding a 221 amino acid ORF that was cloned from a human epithelial cell induced with TNFa and INF cDNA library. The 5' end of HAIDQ59 is represented by the SEQ ID NO: 9, while the 3' end is represented by SEQ ID NO: 10. This clone appears in only two additional cDNA libraries in the HGS database. These two libraries were constructed from the human Jurkat T-cell line and human microvascular endothelial cells. A BLAST analysis of the amino acid sequence of HAIDQ59 demonstrates that this clone exhibits approximately 53% identity and 69% similarity over 226 amino acids of the CD9 TM4SF molecule. (See Figure 8.) It has been demonstrated that the CD9 molecule is involved in signal transduction pathways in platelets, as well as in cell adhesion in both platelets and pre-B-cell lines. Intriguingly, a monoclonal antibody (vpg15), which recognizes the feline homologue of CD9, has been shown to block infection by feline immunodeficiency virus (FIV). Furthermore, a recent study shows that cells expressing high levels of CD9 exhibited suppressed cell motility. Thus, HAIDQ59 may also be involved in signal transduction of blood cells.

The cDNA clone **HHFEK40** contains a 936 nucleotide cDNA insert encoding a 252 amino acid ORF and was cloned from a human fetal heart cDNA library. This clone appears once in the human fetal heart cDNA library and possibly in a hemangiopericytoma cDNA library. A BLAST analysis of the amino acid sequence of .HHFEK40 demonstrated that this clone exhibits approximately 60% identity and 75% similarity over the entire length of a molecule designated PETA-3. (See Figure 9.) PETA-3 was originally identified as a novel human platelet surface glycoprotein termed gp27. Although PETA-3 is present in low abundance on the platelet surface, an anti-PETA-3 monoclonal antibody can stimulate platelet aggregation and mediator release.

WO 98/31799 18 PCT/US98/00959

the transfer of the state of th

Thus, HHFEK40 may function similar to PETA-3 to affect growth of blood cells. Administering polypeptides or fragments of HHFEK40 may be an effective treatment of blood disorders.

The cDNA clone HGBGV89 contains a 738 nucleotide cDNA insert encoding a 197 amino acid ORF and was cloned from a human gall bladder cDNA library. The only two additional appearances of this clone in the HGS database are in a normalized fetal liver cDNA library and in a fetal liver/spleen cDNA library. The cDNA clone HUVBB80 contains a 1071 nucleotide cDNA insert encoding a 201 amino acid ORF and was cloned from a human umbilical vein cDNA library. This clone appears in several additional cDNA libraries in the HGS database including prostate BPH, thyroid, and fetal liver/spleen. BLAST analyses of the amino acid sequences of HGBGV89 and HUVBB80 demonstrate that these clones exhibit approximately 49% identity and 65% similarity and 47% identity and 68% similarity, respectively, over the entire length of a molecule designated L6 surface protein or human tumor-associated antigen L6. (See Figures 10 & 11.) Moreover, another group has confirmed the TM4SF receptor homology of HGBGV89 by describing the protein as a putative transmembrane protein L6H. (See Genbank Accession No 2587054; see Figure 10.) The L6 cell surface antigen is highly expressed on lung, breast, colon, and ovarian carcinomas. Promising results of phase I clinical studies have been reported with an anti-L6 monoclonal antibody, or its humanized counterpart, suggesting that the L6 antigen may be an attractive target for monoclonal antibody-based cancer therapy.

In summary, there is a clear need for identifying and exploiting novel members of the TM4SF superfamily such as those described herein. Although structurally related, these factors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules, such as the novel potential members of the TM4SF superfamily detailed here, should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such factors may prove useful as therapeutics in an anti-tumor, diagnostic, or other capacity. Furthermore, factors such as the nine novel TM4SF superfamily-like molecules described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

Besides TM4SF receptors, receptors from other families are also described. For example, clone **HJACE54**, also called galectin 11, exhibits significant sequence identity to the rategalectin 5, the chicken galectin 3 gene, and the human galectin 8 genes. (See Figure 12.) The galectin 11 cDNA clone contains an 865 nucleotide insert which encodes a 133 amino acid ORF. The clone was obtained from a Jurkat T-cell G1 phase cDNA library. A BLAST analysis of the deduced amino acid sequence of HJACE54 demonstrates approximately 35% identity and 57% similarity to the amino

5

10

15

20

25

30

WO 98/31799 19 PCT/US98/00959

5

10

15

20

25

30

acid sequence of the rat galectin 5 gene. Expression of galectin 11 is quite limited in the HGS database. In fact, the only two additional ESTs in the HGS database which contain the HJACE54 sequence were found in human neutrophil and human infant adrenal gland cDNA libraries. Northern blot analyses have not been performed to examine expression patterns of the galectin 11 gene.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin l appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1mediated disruption of laminin-integrin a₂b₁, interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bcl2, respectively. As a result, the discovery of a novel galectin, such as that encoded by HJACE54, is likely to be a valuable asset both diagnostically and therapeutically.

Additionally, a full-length nucleotide sequence of a novel human cDNA clone which encodes an apparent splice variant of the previously described human E48 antigen has recently been determined. (See Figure 13.) Clone HROAD63 contains a 441 nucleotide cDNA which encodes a 70 amino acid polypeptide. This novel clone exhibits significant sequence identity to several members of a relatively new family of cell-surface proteins termed the Ly6 superfamily. These members include murine and human SCA-2, rat Ly-6 (also termed ThB), and human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)]. The novel E48 splice variant was obtained from the HGS human stomach cDNA library. The clone is present in only a limited number of other HGS cDNA libraries including kidney cancer, keratinocyte, and tongue. An alignment of the nucleotide sequences of the human E48 and HROAD63 cDNAs demonstrates that the initial 168 and 178 nucleotides of E48 and HROAD63, respectively, are identical, with the exception of an additional 10 nucleotides of sequence at the extreme 5' end of the HROAD63 sequence. The

WO 98/31799 20 PCT/US98/00959

Commission Than a promoperation of elegand a productive commission of

5

10

15

20

25

30

35

sequence of the two clones is also identical for an additional 229 nucleotides including the 3' end of the coding sequences and the entire 3' untranslated regions. The only divergence of nucleotide sequence in this region of the clones is the deletion of a single thymidine residue in the 3' UTR of the E48 cDNA. The major difference between the two nucleotide sequences is a 329 nucleotide deletion from the HROAD63 sequence. This deletion causes a shift in the HROAD63 reading frame and encompasses the translational stop signal used in the E48 clone. As a result, the carboxy terminal sequence of HROAD63 is radically altered with regard to that of E48 (as illustrated in Figure 13 by the obvious differences between amino acids 56-128 of E48 and 56-70 of HROAD63 in the amino acid alignment). The clinical presentation of disorders, including abnormal skin and hair phenotypes, may be attributed, at least in part, to a non-functional Ly6 superfamily member such as E48 or HROAD63. HROAD63 may also be involved in blood disorders, as seen with its homologues SCA-2 and CD59.

A novel prohibitin cDNA clone presented herein was originally identified in a human bone marrow cell line (RS4;11) cDNA library. The clone contains a 1066 nucleotide insert which encodes a 299 amino acid polypeptide. BLAST and BestFit analyses of the predicted amino acid sequence of HMWGS46 demonstrate a highly significant sequence identity to a murine protein termed IgM B-cell receptor associated protein (BAP)-37 (Genbank accession number X78683). The HMWGS46 amino acid sequence exhibits nearly perfect identity and similarity over the entire length of the murine BAP-37 sequence. (See Figure 14.) In addition, the full-length nucleotide sequences of HMWGS46 and BAP-37 exhibit at least 87% identical. The HMWGS46 clone also exhibits approximately 49% sequence identity and 85% sequence similarity to a human gene designated prohibitin. Finally, the HMWGS46 cDNA appears in a substantial number of HGS human cDNA libraries in addition to the bone marrow cell line cDNA library from which it was cloned. Some of the cDNA libraries in which this clone appears include keratinocytes, induced endothelial cells, activated neutrophils, synovial sarcoma, colon carcinoma cell line, Jurkat cell line membrane bound polysomes, epileptic frontal cortex, primary dendritic cells, and a number of others. The novel gene related to prohibitin and BAP-37 may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event. Thus, although the precise functional role of the prohibitin family members are less than clear, it is quite likely that such homologues are involved in such complex processes as development, senescence, and tumor suppression. Therefore a novel gene, such as HMWGS46, may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event.

A human cDNA clone encoding a novel epidermal growth factor receptor (EGFR)-like molecule is also disclosed. The novel EGFR-like cDNA clone presented herein was originally identified in an activated human neutrophil cDNA library. The

WO 98/31799 21 PCT/US98/00959

clone contains a 704 nucleotide insert which encodes a 168 amino acid polypeptide. A BLAST analysis of the predicted amino acid sequence of HNFGW06 demonstrates that this novel clone exhibits approximately 85% identity and 90% similarity to a protein designated epidermal growth factor receptor-related protein [Homo sapiens]. (See Figure 15.) The expression profile of the HNFGW06 clone in the HGS database indicates the existence of a fairly highly restricted expression pattern. In addition to the activated neutrophil library from which this clone was obtained, it also appears in the following HGS human cDNA libraries: synovial sarcoma, smooth muscle, placenta, and possibly primary dendritic cells.

The novel EGFR-like cDNA clone HNFGW06 may lead to a number of exciting possibilities for therapeutic and/or diagnostic treatments or reagents. For example, HNFGW06 may be involved in the onset of human breast cancers as well. In addition, due to the fact that TGF-a acts through binding to the EGFR, it is possible that HNFGW06 may also play a role in a variety of gastric processes including regulation of acid secretion, regulation of mucous cell growth, and protection against ethanol- and aspirin-induced injury to gastric tissues.

GENERATING POLYNUCLEOTIDES

5

10

15

20

25

30

35

Polynucleotides of the present invention encoding a receptor may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in cells specified in Table 1 using the expressed sequence tag (EST) analysis (Adams, M.D., et al. Science (1991) 252:1651-1656; Adams, M.D. et al., Nature, (1992) 355:632-634; Adams, M.D., et al., Nature (1995) 377 Supp:3-174.) Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

The nucleotide sequence encoding a receptor polypeptide of SEQ ID NO:Y may be identical to the polynucleotide encoding SEQ ID NO:Y, or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:Y.

When the polynucleotides of the invention are used for the recombinant production of a receptor polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in

Gentz et al., Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Further preferred embodiments are polynucleotides encoding receptor variants comprising the amino acid sequence of receptor polypeptide of Table 1 (SEQ ID NO:Y) in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination.

The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:X or a fragment thereof, or to the cDNA insert in the plasmid deposited at the ATCC, or a fragment thereof, may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding the receptor and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs) that have a high sequence similarity to the receptor gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

In one embodiment, to obtain a polynucleotide encoding the receptor polypeptide, including homologs and orthologs from other species, comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO:X or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to those of skill in the art. Stringent hybridization conditions are as defined above or, alternatively, conditions under overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

5

10

15

20

25

30

WO 98/31799 23 PCT/US98/00959

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

5 Vectors, Host Cells, Expression

10

15

20

25

30

35

The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, and host cells which are genetically engineered with vectors of the invention and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., BASIC METHODS IN MOLECULAR BIOLOGY (1986) and Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, E. coli, Streptomyces and Bacillus subtilis cells; fungal cells, such as yeast cells and Aspergillus cells; insect cells such as Drosophila S2 and Spodoptera Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., MOLECULAR CLONING, A LABORATORY MANUAL (supra).

24 WO 98/31799 PCT/US98/00959

· Som it is to the plant of a consequence of

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If the receptor polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the receptor polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Receptor polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

20 Diagnostic Assays

5

10

15

25

30

This invention also relates to the use of receptor polynucleotides or polypeptides for use as diagnostic reagents. Detection of a mutated form of the receptor gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a disease or susceptibility to a disease which results from underexpression, over-expression or altered expression of the receptor. Individuals carrying mutations in the receptor gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled receptor nucleotide sequences. Perfectly matched 35 sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. (See, e.g., Myers et al., Science (1985) 230:1242.) Sequence changes at specific locations may also be revealed by

WO 98/31799 25 PCT/US98/00959

nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method. (See Cotton et al., Proc Natl Acad Sci USA (1985) 85: 4397-4401.) In another embodiment, an array of oligonucleotides probes comprising receptor nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability. (See for example: M.Chee et al., Science, Vol 274, pp 610-613 (1996).)

The diagnostic assays offer a process for diagnosing or determining a susceptibility to specific diseases through detection of mutation in the receptor gene by the methods described.

In addition, specific diseases can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of receptor polypeptide or receptor mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease which comprises:

- (a) a receptor polynucleotide, preferably the nucleotide sequence of SEQ ID NO:X, or a fragment thereof;
 - (b) a nucleotide sequence complementary to that of (a);
- (c) a receptor polypeptide, preferably the polypeptide of SEQ ID NO:Y, or a fragment thereof; or
- (d) an antibody to a receptor polypeptide, preferably to the polypeptide of SEQ ID NO: Y.
- It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Chromosome Assays

5

10

15

20

25

The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the

WO 98/31799 26 PCT/US98/00959

sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

10

15

20

25

30

35

5

Antibodies

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the receptor polypeptides. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against the receptor polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., Nature (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., Immunology Today (1983) 4:72) and the EBV-hybridoma technique (Cole et al., MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against receptor polypeptides may also be employed to treat diseases.

Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with a receptor polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immune response to protect said animal from a disease. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which

WO 98/31799 27 PCT/US98/00959

comprises, delivering a receptor polypeptide via a vector directing expression of the receptor polynucleotide in vivo in order to induce such an immunological response to produce antibody to protect said animal from diseases.

Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a receptor polypeptide wherein the composition comprises a receptor polypeptide or receptor gene. The vaccine formulation may further comprise a suitable carrier. Since a receptor polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation instonic with the blood of the recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

Screening Assays

5

10

15

20

25

30

35

The receptor polypeptide of the present invention may be employed in a screening process for compounds which bind the receptor and which activate (agonists) or inhibit activation of (antagonists) the receptor polypeptide of the present invention. Thus, polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).

The receptor polypeptides are responsible for many biological functions, including many pathologies. Accordingly, it is desirous to find compounds and drugs which stimulate the receptor on the one hand and which can inhibit the function of the receptor on the other hand. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions and diseases. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions and diseases.

In general, such screening procedures involve producing appropriate cells which express the receptor polypeptide of the present invention on the surface thereof.

10

15

20

25

30

Such cells include cells from mammals, yeast, Drosophila or E. coli. Cells expressing the receptor (or cell membrane containing the expressed receptor) are then contacted with a test compound to observe binding, or stimulation or inhibition of a functional response.

The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the receptor is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the receptor, using detection systems appropriate to the cells bearing the receptor at their surfaces. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

Further, the assays may simply comprise the steps of mixing a candidate compound with a solution containing a receptor polypeptide to form a mixture, measuring receptor activity in the mixture, and comparing the receptor activity of the mixture to a standard.

The receptor cDNA, protein and antibodies to the protein may also be used to configure assays for detecting the effect of added compounds on the production of receptor mRNA and protein in cells. For example, an ELISA may be constructed for measuring secreted or cell associated levels of receptor protein using monoclonal and polyclonal antibodies by standard methods known in the art, and this can be used to discover agents which may inhibit or enhance the production of the receptor (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues. Standard methods for conducting screening assays are well understood in the art.

Examples of potential receptor antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligand of the receptor, e.g., a fragment of the ligand, or small molecules which bind to the receptor but do not elicit a response, so that the activity of the receptor is prevented.

Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for receptor polypeptides; or compounds which decrease or enhance the production of receptor, which comprises:

- (a) a receptor polypeptide, preferably that of SEQ ID NO:Y;
- (b) a recombinant cell expressing a receptor polypeptide, preferably that of SEQ 35 ID NO:Y;
 - (c) a cell membrane expressing a receptor polypeptide; preferably that of SEQ ID NO: Y; or
 - (d) antibody to a receptor polypeptide, preferably that of SEQ ID NO: Y.

SUBSTITUTE SHEET (RULE 26)

WO 98/31799 29 PCT/US98/00959

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Prophylactic and Therapeutic Methods

5

10

15

20

25

30

35

This invention provides methods of treating an abnormal conditions related to both an excess of and insufficient amounts of receptor activity.

If the activity of the receptor is in excess, several approaches are available. One approach comprises administering to a subject an inhibitor compound (antagonist) as described along with a pharmaceutically acceptable carrier in an amount effective to inhibit activation by blocking the binding of ligands to the receptor or by inhibiting a second signal, and thereby alleviating the abnormal condition.

In another approach, soluble forms of the receptor polypeptides still capable of binding the ligand in competition with endogenous receptor may be administered. Typical embodiments of such competitors comprise fragments of the receptor polypeptide.

In still another approach, expression of the gene encoding endogenous receptor can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. (See, for example, O'Connor, J Neurochem (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Alternatively, oligonucleotides which form triple helices with the gene can be supplied. (See, for example, Lee et al., Nucleic Acids Res (1979) 6:3073; Cooney et al., Science (1988) 241:456; Dervan et al., Science (1991) 251:1360.) These oligomers can be administered per se or the relevant oligomers can be expressed in vivo.

For treating abnormal conditions related to an under-expression of the receptor and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates the receptor, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of the receptor by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells in vivo and expression of the polypeptide in vivo. For overview of gene therapy, see Chapter 20, Gene Therapy and

15

20

25

30

35

- Contract of the Calegory distribution of weather that the latter than the calegory of the calegory of the calegory

other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996).

5 Formulation and Administration

Peptides, such as the soluble form of receptor polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds. Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels and the like.

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of $0.1\text{-}100\,\mu\text{g/kg}$ of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide ex vivo, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

Paramana kalabar kabupat kalendar kelabah Di Maralah Kerada Kalabah Kerada Kalabah Kerada Kaban Kerada Kaban K

SUBSTITUTE SHEET (RULE 26)

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

SUBSTITUTE SHEET (RULE 26)

Page 1

	HMACR70																													
_			Ţ										+									AGAT	+				<u> </u>		+	90
ĊGT	CGTCAAGGACTCTCTTCTTGGGACTCCTTGTCTGCAAGGGAGCGCCGGGGACCGTGGAGGTTGGGGTCTATACGACGACGACGACGACGACGAC														,															
																							M	L	L	L	L	L	L	
ccc	CT.	GCTC	TGG	GG	SAGG	GAG	AGG	GTG	GA.	AGGA	CAG	AAG	AG	ΓΑΑ	CGG	AAG	GAT	TAC	TC	сто	AC	GATG	CAG	AGT	TC	CGTO	SACC	GTG	CAA	180
GGG	GA	CGAG	ACC	CCC	CTCC	cto	TCC	CAC	СТ	тсст	GTC	TTC	TC	ATTO	GCC	TTC	CTA	ATG	AGC	GAC	TG	CTAC	GTC	TCA	AG	GCAC	TGG	CAC	GTT	,00
Ρ	L	L	W	G	R	Ε	R	٧	Ε	G	۵	κ	s	N	R	K	0	Υ	s	L	Т	M	Q	s	s	ν	т	ν	a	
GAG	GG	CATG	TGT	GTO	CAT	rgtg	CGC	TGC	TC	CTTC	тсс	TAC	ÇC/	GTO	GAC	AGC	CAG	ACT	GAC	TCT	GA	CCCA	GTT	CAT	GG	CTĄ	CTGG	TTC	ceė	070
сто	CC	GTAC	ACA	CAC	GTA	CAC	GCG	ACG	AG	GAAG	AGG	ATG	GG	CAC	ĊTG	TCG	GTC	TGA	CTO	SAGA	C T	GGGT	CAA	GTA	CC	GAT	SACC	AAG	GCC	270
E	G	м	С	٧	н	٧	R	С	s	F	s	Y	P	٧	D	s	0	τ	D	s	D	Ρ	٧	н	G	۲	W	F	R	
GCA	GG	GAAT	GAT	ΆΤΑ	AGC	TGG	AAG	GCT	CC.	AGTG	GCC	ACA	ΑΑ	CAAC	CCA	GCT	TGG	GCA	GTG	CAG	GA	GGAA	ACT	CGG	GAI	CCGA	ATTC	CAC	CTÇ	
CGT	cc	CTTA	CTA	TAI	TCC	ACC	TTC	CGA	GG	TCAC	CGG	TGT	TT	STTO	GGT	CGA	ACC	CGT	CAC	GTO	CT	CCTT	TGA	GCC	CT	GGC	ΓAAG	GTG	GAG	360
Α	G	N	O	ì	5	W	к	A	P	٧	Α	τ	N	N	P	A	٧	А	٧	٥	Ε	Ε	τ	R	D	R	F	н	L	
CTI	GG	GGAC	CCA	CAC	SACO	CAAA	AAT	TGC	AC	CCTG	SAGO	ATC	AGA	GAI	recc	AGA	ATG	SAGT	GAT	GCG	GG	GAGA	TAC	TTC	TT	TCG	FATG	GAG	AAA	
GAA	CC	CCTG	GGT	GTO	TGG	 	TTA	ACG	TG	GGAC	TCG	TAG	TCI	CTA	CGG	TCT	TAC	TCA	CTA	CGC	CC	стст	ATG	AAG	AA	AGC	TAC	сто		450
L	G	D	P	Q	T	κ	N	С	т	L	s	1	R	D	A	R	M	s	D	A	G	R	Y	F	F	R	м	Ε	ĸ	
GGA	AA	TATA	AAA	TGC	SAAT	TAT	AAA	TAT	GA	CCAG	стс	тст	GTO	SAAC	GTG	ACA	TAC	CCT	CCI	CAG	AA	CTTG	ACT	GTG	AC.	TGTO	TTC	CAA	GGA	
CCT	TT.	ATAT	777	ACC	TTA	ATA	TTT	ATA	CT	GGTC	GAG	AGA	CAC	TTO	CAC	TGT	ATO	GGA	GGA	GTO	TT	GAAC	TGA	CAC	TG	ACAC	SAAG	GTT	ССТ	540
G	N	1	к	W	N	۲	κ	Y	Đ	a	L	s	٧	N	٧	T	Y	ρ	Ρ	a	N	L	τ	V	T	٧	F	a	G	
GAA	(GG)	CACA	GCA	TCC	CACA	AGCT	CTG	GGG	AAI	CAGO	TCA	TCT	CTI	rtc	AGTC	СТА	GAG	seec	CAG	STCT	CT	GCGC	TTG	GTC	TG	TGCI	GTT	GAC	AGC	
			-							+			+									CGCG	+							630
																						R								
AAT	CC	CCC1	GCC	AGO	CTG	SAGC	TGG	ACC	TG	GAGG	AGT	CTG	AC (сто	TAC	ccc	TCA	CAG	ccc	TÇA	AAI	CCCT	CTG	GTA	ст	GGAC	CTG	CAA	GTG	
	-+		+			-+-		+		+			+			-		-				GGGA	+					—		720
N	Р	P	Α	R	L	s	W	T	W	R	s	L	Т	L	Y	ρ	s	a	Ρ	s	N	Р	L	٧	L	Ε	L	a	v	
CAC	CT.	GGGG	GAT	GA/	.GGG	GAA	TTC	ACC	TG	TCGA	GCT	CAG	AA(:TC1	rcts	GGT	TCC	CAG	CAC	GTT	TC	CCTG	AAC	стс	TC	сто	CAA	CAG	GAĢ	010
GTG	GA	CECC	CTA	CTI	CCC	CTT	AAG	TGG	AC.	AGCT	CGA	GTC	TTO	SAGA	GAC	CCA	AGG	BTC	GTO	CAA	AG	GGAC	TTG	GAG	AG	GGAC	GTT	GTC	CTC	810
н	L	G	Đ	Ε	G	Ε	F	Т	С	R	Α	a	N	s	L	G	s	Q	н	٧	s	Ĺ	N	L	s	L	a	a	Ε	

an tark flank the flank the stark the second the second the second the second the second that the second the s

Page

		HTEDK48 CGAGTGGACAACCATCAGGGGGGGGGGGGGGGGGGGGG																												
ĆĠ	GC A	CGA	GŢG	SACA	AACC	CATO	CAGO	GAC	SCC/	AGGA	CAC	AGA	GGC	GC/	GAG	CAA	GTO	AGC	ATT	GGC	GCC	CCT	ŢCC	TC	GAT	ccc	TAT	CAT	CTT	
GC	GT	GCT	CAC	TG1	TTG	STAG	STCC	CTC	GGT	TCCI	GTG	TCT	ccc	CGI	стс	GTT	CAG	TCG	TAA	CCG	CGG	GGA	AGG	AG 1	CTA	GGG	ATA	GTA	GAA	90
	GGGAAACAGTAGCCCAGAGGTTCAGGAAGATGTTAACTTAAATGTTCGGGGTGCCCCAGTCTGTTCAGCATGGCTGAAATCCACACTCC																180													
CC	TT.	TGT	CAT	:GGC	STCT	CCA	I A G T	CCT	TCI	ΓΑΟΑ	ATT	GAA	TTI	ACA	AGC	ccc	A C G	GGG	TCA	GAC	AAG	TCG	TAC	CGA	CTT	TAG	GTG	TGA	GGC	
																							н	A	Ε	ł	Н	T	P	
TAT	TC	TTC	TTC	SAAG	SAAA	CTG	STTA	TCT	TTA	ACTO	AAT	GGC	TTC	GTG	GCT	GTG	TCT	GGC	ATÇ	ATC	CTA	GTT	GGC	CTG	GGC	ATT	GGT	GGT	AAA	
ATA	AGA	AAG	SAAC	:110	777	GAC	CAAT	AGA	AAT	GAG	TTA	CCG	AAG	CAC	CGA	CAC	AGA	CCG	TAG	TAG	GAT	CAA	ccs	GAC	CCG	TAA	CCA	CCA	TTT	270
Y	s	s	L	к	K	L	L	S	L	L	N	G	F	٧	A	٧	s	G	Ī	I	L	٧	G	L	G	į	G	G	к	
TG1	GGA	4G G (igco	тст	гстс	ACG	SAAT	GTC	сто	GGG	CTG	TCC	ŢCC	GCA	TAC	CŢC	стт	CAC	GTT	GGC	AAC	CTG	TGC	CTG	GTG	ATG	GGA	TGC	ATC	
ACA	CC	rcco	CGG	AGA	GAC	TGC	TTA	CAG	GAG	ccc	GAC	AGG	AGG	CGT	ATG	GAG	GAA	G TG	CAA	CCG	T TG	GAC	ACG	GAC	CAC	TAC	ССТ	ACG	TAG	360
С	G	G	A	S	Ĺ	Т	N	٧	Ł	G	L	s	s	A	Υ	L	L	н	٧	G	N	L	ε	L	٧	M	G	С	ī	
ACG	GTA	СТО	ÇTT	GGC	TGT	GCC	GGG	TGG	TAT	GGA	GCG	AC T	AAA	GAG	AGC	AĢA	GGC	ACG	СТС	TTG	TTT	TGC	ATC	CTG	TCA	ATG	GTT	A T T	GTÇ	
TGC	CAT	GAC	GAA	CCG	ACA	CGG	ccc	ACC	ATA	CCT	CGC	TGA	TTT	СТС	TCG	TCT	CCG	TGC	GAG	AAC	AAA	ACG	TAG	GAC	AGT	TAC	CAA	TAAI	CAG	450
T	ν	L	L	G	С	A	G	W	Υ	G	A	T	ĸ	Ε	s	R	G	T	L	L	F	С	Į	L	s	м	γ	ı	V	
СТС	ATC	ATG	GAA	GTT	ACA	GCT	GCC	ACA	GTG	GTC	CTT	CTT	TTC	T T T	CCA	ATT	GTT	GGA	GAT	GTG	GCC	TTG	GAA	CAC	ACC	TTC	GTG	ACCI	CTG	
GAG	TAG	TAC	CTT	CAA	TGT	CGA	CG G	TGT	CAC	CAG	GAA	GAA	AAG	AAA	GGT	TAA	CAA	ССТ	CTA	CAC	cgg	AAC	CTT	GTG	TGG	AAG	CAC	TGG	GAC	540
L	1	M	Ε	٧	Т	A	A	т	V	V	L	L	F	F	P	i	٧	G	0	٧	A	L	Ξ	н	Т	F	٧	Т	L	
AGG	AAG	AAT	ŢAC	AGA	GGT	TAC	AAC	GAG	CCA	GAC	GAC	TAT	тст	ACA	CAG	TĢG.	AAC	TTG	G TÇ.	4 T G i	GAG.	AAG	ÇTA	AAG	TGC	TGT	GGG	STG/	AAT	
TCC	TTC	TTA	ATG	TCT	CCA.	ATG	TTG	стс	GGT	CTG	C TG	ATA,	AGA	TGT	GTC.	ACC	TTG	AAC	CAG	TAC	стс	TTC	GAT	TTC	A CG	ACA	ccc	CACT	TTA	630 ~
R	к	N	Y	R	G	Y	N	Ε	P	О	a	Y	s	T	۵	W	N	L	٧	M	Ε	κ	L	κ	С	С	G	ν	N	
AAC	TAC	ACA	ĢAT	TTT	TCT	GGC.	TCT	TCC	T T Ç	GAA.	A TG	ACA.	ACG	GGC	CAC	AČC.	TAC	CC C /	AGGA	AGT.	TGC	TGT	AAA'	TCC.	ATC	GGA	AGTO	STGT		
ŤTG	ATG	TGT	CTA	AAA.	AGA	CCG	AGA,	AG G	AAG	CTT	TAC	TGT	TGC	CCG	GTG	T G G.	A TG	GGG	TCC	ГСА	ACG.	ACA	177,	4GG	TAG	CCT	CAC	CACA		720
N	Y	Т	О	F	s	G	s	s	F	Ε	M	T	т	G	н	т	Y	P	R	s	С	С	к	s	Į	G	s	ν	s	
TGT	GAC	GGA	ÇGC	GAT	GTG:	TÇTI	CCA	AACI	GTC.	ATC	CA C (CAG	4 A G	GGC	TGT	TTC	CAT	AAAC	стсс	TA	AAA	ATC	ACC/	٩AG	AC T	CAGA	AGC1	TCA		
ACA	CTG	CCT	GCG	CTA	CACA	AGA	GGT.	TTG	CAG	TAG	GTG	STC	TTC	ccG,	ACA/	4ÅG(GTA	TTTC	SAGO	AT	TT	TAG	TGG	TC	TGA	STC	CGA	AGT	rg G	810
С	Ð	G	R	D	٧	s	P	N	v	Į	н	Q	к	G	С	F	Н	ĸ	L	L	ĸ	Į	7	ĸ	т	Q	s	F	T	
CTG.	AGT	GGG	AGC.	TCT	CTG	GGA	GCT(GCA	STG.	ΑΤΑΙ	CAGA	GG	r G G (GGG.	тста	GC:	TATO	STTO	scgo	AGO	CTO	GTO	:110	SAAI	CTG	CTGO	CCT	'AAA		GID NO:3
GAC	TCA	ССС	TCG	AGA	GAC	ССТ	CGAC	CGT	CAC	TATO	STCT	CC/	CC	CCC	AGAC	G C G	ATA	CAAC	GCG	TC	GA	CAC	SAAC	TTC	SACO	SACO	GGA	TTT	CG	900
L	s	G	5	S	L	G	A	A	٧	i	a	R	W	G	s	R	Y	٧	A	a	A	G	L	Ε	L	L	A	. رد		ID 19

age 1

HPWA E25	
GGCGTCCCTCTGCCTGCCCACTCAGTGGCAACACCCGGGAGCTGTTTTGTCCTTTGTGGAGCCTCAGCAGTTCCCTCTTTCAGAACTCA	
CCGCAGGGAGACGGACGGGTGAGTCACCGTTGTGGGCCCTCGACAAAACAGGAAACACCTCGGAGTCGTCAAGGGAGAAAGTCTTGAGT	- 90 }
TGCCAAGAGCCCTGAACAGGAGCCACCATGCAGTGCTTCAGCTTCATTAAGACCATGATGATCCTCTTCAATTTGCTCATCTTTCTGTG	,
ACGGTTCTCGGGACTTGTCCTCGGTGGTACGTCACGAAGTCGAAGTCATCTGGTACTACTAGGAGAAGTTAAACGAGTAGAAAGACAC	. 180
MUCFSFIKIMMILFNEEIFEC	
GGTGCAGCCCTGTTGGCAGTGGGCATCTGGGTGTCAATCGATGGGGCCATCCTTTCTGAAGATCTTCGGGCCACTGTCGTCCAGTGCCAT	- 270
CCACGTCGGGACAACCGTCACCCGTAGACCCACAGTTAGCTACCCCGTAGGAAAGACTTCTAGAAGCCCGGTGACAGGTCACGGTA	
G A A L L A V G I W V S I D G A S F L K I F G P L S S S A M	
CAGTTTGTCAACGTGGGCTACTTCCTCATCGCAGCCGGCGTTGTGGTCTTTGCTCTTGGTTTCCTGGGCTGCT	:
GTCAAACAGTTGCACCCGATGAAGGAGTAGCGTCGGCCGCAACACCAGAAACGAGAACCAAAGGACCCGACGA	- 360
GTCAAACAGTTGCACCCGATGAAGGAGTAGCGTCGGCCGCAACACCAACAACAAAGGACCCGACGATACCACGATTCTGACT	
Q F V N V G Y F L I A A G V V V F A L G F L G C Y G A K T E	
AGCAAGTGTGCCCTCGTGACGTTCTTCTTCATCCTCCTCCTCATCTTCATTGCTGAGGTTGCAGCTGCTGTGGTCGCCTTGGTGTACAC	
TCGTTCACACGGGAGCACTGCAAGAAGAAGTAGGAGGAGGAGTAGAAGTAACGACTCCAACGTCGACGACACCAGCGGAACCACATGTG	- 450 :
ACAATGGCTGAGCACTTCCTGACGTTGCTGGTAGTGCCTGCC	- 540
TGTTACCGACTCGTGAAGGACTGCAACGACCATCACGGACGG	
T.M.A.E.H.F.L.T.L.V.V.P.A.I.K.D.Y.G.S.Q.E.D.F.T.Q.V.W.N	
ACCACCA TGAAAGGGC TCAAGTGC TGTGGC TTCACCAACTA TACGGA TTTTGAGGAC TCACCC TACTTCAAAGAGAACAGTGCC TTTCC	
TGGTGGTACTTTCCCGAGTTCACGACACCGAAGTGGTTGATATGCCTAAAACTCCTGAGTGGGATGAAGTTTCTCTTGTCACGGAAAGG	630
Tad fact free conditions and tad fact from the condition of the condition of the conditions and the conditions and the conditions are conditional to the conditions and the conditions are conditional to the conditional to the conditions are conditional to the condit	
TTMKGLKCCGFTNYTDFEDSPYFKENSAFP	
CCATTCTGTTGCAATGACAACGTCACCAACACAGCCAATGAAACCTGCACCAAGCAAAAAGGCTCACGACCAAAAAGTAGAGGGTTGCTT	720
GGTAAGACAACGTTACTGTTGCAGTGGTTGTGTCGGTTACTTTGGACGTGGTTCGTTTTCCGAGTGCTGGTTTTTCATCTCCCAACGAA	720
P F C C N D N V T N T A N F T C T K D K A H D D K V E G C F	
AATCAGCTTTTGTATGACATCCGAACTAATGCAGTCACCGTGGGTGG	810
TTAGTCGAAAACATACTGTAGGCTTGATTACGTCAGTGGCACCCACC	
N Q L L Y D I R T N A V T V G G V A A G I G G L E L A A M N	
TGTGTCCATGTATCTGTACTGCAATCTACAATAAGTCCACTTCTGCCTCTGCCACTACTGCTGCCACATGGGAACTGTGAAGAGGCACC	
ACACAGGTACATAGACATGACGTTAGATGTTATTCAGGTGAAGACGGAGACGGTGATGACGGTGTACCCTTGACACTTCTCCGTGG	900
TO THE TOTAL CONTROL OF THE TOTAL CONTROL OF THE TOTAL CONTROL OF THE CONTROL OF	

HPWAF25

Page 2

TGGCAAGCAGCAGTGATTGGGGGAGGGGACAGGATCTAACAATGTCACTTGGGCCAGAATGGACCTGCCCTTTCTGCTCCAGACTTGGGG	990
ACCGTTCGTCGTCACTAACCCCCTGTCCTAGATTGTTACAGTGAACCCGGTCTTACCTGGACGGGAAAGACGAGGTCTGAACCCC	330
WOAAVIGGGDRI. (SOO, IDNO! 20)	
CTAGATAGGGACCACTCCTTTTAGGCGATGCCTGACTTTCCTTCC	1080
GATCTATCCCTGGTGAGGAAAATCCGCTACGGACTGAAAGGAAGG	1000
GCCAGTTCTGTTGCCCATTCCCCCAGTCTATTAAACCCTTGATATGCCCCCTAGGCCTAGTGGTGATCCCAGTGCTCTACTGGGGGATGA	1170
CGGTCAAGACAACGGGTAAGGGGGTCAGATAATTTGGGAACTATACGGGGGATCCGGATCACCACTAGGGTCACGAGATGACCCCCTACT	1170
GAGAAAGGCATTT-TATAGCCTGGGCATAAGTGAAATCAGCAGAGCCTCTGGGTGGATGTGTAGAAGGCACFTCAAAATGCATAAACCTGT	
	1260
CTCTTTCCGTAAAATATCGGACCCGTATTCACTTTAGTCGTCTCGGAGACCCACCTACACATCTTCCGTGAAGTTTTACGTATTTGGACA	



38

Päge 1

	HTPEF86																															
AAA	AA	AAA	CAA	GG	TCC	CCA	CA	GCA	AAC	AA	AAGG	AA	TAG	GA T	CAA	GAG	ATA	ACGT	GGC	TGC	TG	GCA	GA	GCA	AGC						TTCA	
TTI	ΤI	TTT	GTT	CC	AGG	GGT	GT	CGT	TTO	TT	TTCC	TTA	TC	CTA	GTT	TC	TA	TGC	CCG	ACC	AC	CGT	CT	CGT	TCG	TAC	TT.	AAG	CTAC	:TG	AAGT	90
																										M	N	S	M	T	, \$	•
GCA	GT	T C C	GĢT	GG	CÇA.	ATT	СT	GTG	TTO	GT	GGTG	GC/	AC C (CÇA	CAA	TGG	TTA	ATEC	TGT	GAC	ĊC	CAG	GA	ATT	ATG	TCI	CA	CGT	3000	СТ	GTAŢ	
CGI	C.A	AGG	CCA	cc	GGT	TAA	GA	CAC	AAC	CA	CCAC	CG1	GG	GT	GTT	ACC.	AA1	rago	ACA	CTO	GG	GTC	CT	TAA	TAC	AGA	GT	GCA	GGG	G A	CATA	180
A	٧	<i>i</i> P	ν		A	N	S	٧	L	γ	ν	A	P	н	N	G	١	Y F	, Λ	1	r	P	G	1	Ħ	S	н	ν	P	L	Y	
CCA	AA	CAG	CÇA	GC	CGC	AAG	TC	CAC	CTA	AGT	гсст	GGC	AA	çç.	ACC1	rag:	ΓŦŢ	rggi	GTC	GAA	ĄŢG	TGA	AT	GGG	ÇAG	C C 1	GT	GCĄ	SAAA	rec.	TCTG	
GGT	ΤĪ	GTC	GGT	CG	GCG	TTC	AG	GTG	GA 1	C A	AGGA	CCC	TTO	GG	TGG	ATC.	444	ACC.	CAG	CTT	TAC	ACT	TA	ccc	GTC	GGA	CA	CGT	: T T T	CG.	AGAC	270
Ρ	N	ı s	a		P	a	٧	н	L	٧	P	G	N	Р	Р	s	Ł	٠ ،	s		ı	٧	N	G	Q	P	٧	Q	κ	Α	L	
AAA	GA	AGG	CAA	AA	CCT	TGG	GG	GC C	ATC	CA	GATC	ATO	AT ²	rggi	ССТО	GC.	TCA	LA3	CGG	CCT	reg	GET	cc.	ATC	ATG	GCG	AC	GGT	rctc	GT.	AGGG	
TTT	CT	TCC	GTT	TT	GGA	ACC	CC	CGG	TAG	GT	TAG	TAG	TA	YCC!	GGAC	CGA	\GT	TGTA	GCC	GGA	GC	C GA	GG	TAG	TAC	CGC	TG	CÇA/	GAG	CA	TCCC	360
к	Ε	: G	K		T i	L	G	Α	ł	0	ı	ı	1	G	L	A	۲	4 1	G	L		G	s	ı	M	Α	т	٧	L	٧	G	
GAA	TΑ	CCT	GTC	TΑ	TTT	CAT	TC.	TAC	GGA	.GG		ccc	TT(TG	GGGA	AGGO	: T T	rete	GTT	TAT	·£ A	דדד	CA	GGA	TCT	сто	TC	CGTO	GC.A	AGC.	AGAA	
			- j				+							-							+-				+							450
		, F																											А			
		_							Ī	Ī																						
			- i -											+-							-				+						CTTC	540
TTA	GT.	CGG	TAT	AAI	GAAT	ΓΑΑ	CGC	SAC	GAC	AGA	ACCG	TCA	AA(: C C (SAAC	TTO	T A	AGCA	GTC	ACG	TT	AGA	CG	AGA	CGT	CAA	CC.	rcae	TAT	GA	GAAG	
N	٥	P	Υ	;	s '	Y	С	L	L	S	G	s	L	G	L	N	I	Į V	S	A	١	i	С	s	A	٧	G	٧	i	L	F	
ATC	AC	AGA	TÇT	AA	GTA	TTC	ĊСС	CAC	CCA	TĄ	rgcc	TAC	ccc	GAC	TAT	TAT	rcc	TTA	CGC	CTG	ĢG	GTG	TG	٩AC	сст	GGA	ATO	GÇC	ATT	TC	TGGÇ	620
TAG	TG	TCT	AGA	TT	CATA	AAG	GG	STG	GGT	A T	cee	ATG	GGG	CTO	ATA	ATA	GG	AAT	GCG	GAC	cc	CAC	AC	TTG	GGA	CCT	TAC	ccc	TAA	AG	A C C G	630
I	T	D	L	;	S	ı	٩	Н	P	Y	A	Y	Ρ	D	Y	Y	P	> Y	Α	W	ı	G	٧	N	P	G	M	A	1	s	G	
GT G	СT	GCT	GGT	C T	TCTO	scc.	TCC	TG	GAG	TTI	GGC	ATC	GC A	TG	GCA	TCT	тс	CCA	стт	TGG	C T	GC C	AG:	FTG	GTC	TGC	TGI	ГСАА	TCA	AGO	CAAT	
CAC	GA	CGA	CCA	G A A	AGAC	GG	AGO	SAC	СТС	AAA	CCG	TAG	CG1	ACO	CGT	AGA	۱AG	GGT	GAA	AC C	GA	CGG	TC	AAC	C AG	ACG	AC/	GTT	AGT	TC	STTA	720
٧	L	. L	٧	ę	- (: (L	L	E	F	G	ı	Α	С	Α	s	S	5 Н	F	G	:	С	O	L	٧	С	С	۵	s	s	N	
GT G	AG	TGT	CAT	C T A	ATCO	:AA	AC.	ATC:	TAT	GCA	GCA	AAC	CCA	GTO	SATO	ACC	:cc	AGA	ACC	GGT	GA	ССТ	CAC	CA	CCA	AGT	TAl	rtcc	AGT	GAG	SATC	
			-+-				+							-							+-		-					-				810
٧	s	. V	i	,	Y F	۱ د	N	ſ	Y	Α	Α	N	P	٧	ı	Т	Р	, Е	Р	٧		Т	s	P	P	s	Υ	s	s	Ε	ı	

Päge 1

HSBRF02	
CACGAGCAGGGTCTCGGGCTAGTCATGGCGTCCCCGTCTCGGAGACTGCAGACTAAACCAGTCATTACTTGTTTCAAGAGCGTTCTG	CTA
GTGCTCGTCCCAGAGCCCGATCAGTACCGCAGGGGCAGAGCCTCTGACGTCTGATTTGGTCAGTAATGAACAAAGTTCTCGCAAGAC	GAT
MASPSRRLQTKPVITCFKSVL	۲.
ATCTACACTTTTATTTTCTGGATCACTGGCGTTATCCTTCTTGCAGTTGGCATTTGGGGCAAGGTGAGCCTGGAGAATTACTTTTCT	СТТ
TAGATGTGAAAATAAAAGACCTAGTGACCGCAATAGGAAGAACGTCAACCGTAAACCCCGTTCCACTCGGACCTCTTAATGAAAAGA	 1 180 Gaa
I Y T F I F W I T G V I L L A V G I W G K V S L E N Y F S	L
TTAÁATGAGAAGGCCACCAATGTCCCCTTCGTGCTCATTGCTACTGGTACCGTCATTATTCTTTTGGGCACCTTTGGTTGTTTTGCT	ACC
AATTTACTCTTCCGGTGGTTACAGGGGAAGCACGAGTAACGATGACCATGGCAGTAATAAGAAAACCCGTGGAAACCAACAAAACGA	+ 270 ⊺GG
L N E K A T N V P F V L I A T G T V I I L L G T F G C F A	т
TGCCGAGCTTCTGCATGGATGCTAAAACTGTATGCAATGTTTCTGACTCTCGTTTTTTTGGTCGAACTGGTCGCTGCCATCGTAGGA	TTT
ACGGCTCGAAGACGTACCTACGATTTTGACATACGTTACAAAGACTGAGAGCAAAAAAAA	+ 360 AAA
C R A S A W M L K L Y A M F L T L V F L V E L V A A I V G	F
GTTTTCAGACATGAGATTAAGAACAGCTTTAAGAATAATTATGAGAAGGCTTTGAAGCAGTATAACTCTACAGGAGATTATAGAAGC	CAT
CAAAAGTCTGTACTCTAATTCTTGTCGAAATTCTTATTAATACTCTTCCGAAACTTCGTCATATTGAGATGTCCTCTAATATCTTCG	-+ 450
V F R H E I K N S F K N N Y E K A L K Q Y N S T G D Y R S	н
GCAGTAGACAAGATCCAAAATACGTTGCATTGTTGTGGTGTCACCGATTATAGAGATTGGACAGATACTAATTATTACTCAGAAAAA	3GA
CGTCATCTGTTCTAGGTTTTATGCAACGTAACAACACCACAGTGGCTAATATCTCTAACCTGTCTATGATTAATAATGAGTCTTTTTC	─+ 540 CT
A V D K I Q N T L H C C G V T D Y R D W T D T N Y Y S E K	G
TTTCCTAAGAGTTGCTGTAAACTTGAAGATTGTACTCCACAGAGAGATGCAGACAAAGTAAACAATGAAGGTTGTTTTATAAAGGTG	√TG
AAAGGATTCTCAACGACATTTGAACTTCTAACATGAGGTGTCTCTCTACGTCTGTTTCATTTGTTACTTCCAACAAAATATTTCCACT) 630 ΓΑC
F P K S C C K L E D C T P Q R D A D K V N N E G C F I K V	M
ACCATTATAGAGTCAGAAATGGGAGTCGTTGCAGGAATTTCCTTTGGAGTTGCTTGC	' G C
TGGTAATATCTCAGTCTTTACCCTCAGCAACGTCCTTAAAGGAAACCTCAACGAACG	+ 720 ICG
T I I E S E M G V V A G I S F G V A C F Q L I G I F L A Y	С
CTCTCTCGTGCCATAACAAATAACCAGTATGAGATAGTGTAACCCAATGTATCTGTGGGCCTATTCCTCTACCTTTAAGGACATTT	AG
GA GAGAGCÁC GGTA TIGITTATT GGTCATACT CTATCACATT GGGTT ACATAGAGAC CCGGATAAGGAGAGAT GGAAATT CCTGTAAA	<u></u>
LSRALTNNOYELV. (SCG. JD NO: 22)	
/ GGTCCCCCTGTGAATTAGAAAGTTGCTTGGCTGGAGAACTGACAACACTACTTACT	.TT
	900

Page 1

HLTAHO	
CACGAGCATTGCCGCTCTCTCGGTGAGCGCAGCCCCGCTCTCCGGGGCCTTCGCGGGCCACCGGCGCATGGGCCAGTGCGGCATC	90
GTGCTCGTAACGGCGAGAGACCACTCGCGTCGGGGCGAGAGGCCCGGAAGCGCCCGGTGGCCGCGGTACCCCGGTCACGCCGTAG	90
M G O C G I	
ACCTCCTCCAAGACCGTGCTGGTCTTTCTCAACCTCATCTTCTGGGGGGCAGCTGGCATTTTATGCTATGTGGGAGCCTATGTCTTCATC	180
TGGAGGAGGTTCTGGCACGACCAGAAAGAGTTGGAGTAGAAGACCCCCCGTCGACCGTAAAATACGATACACCCTCGGATACAGAAGTAG	100
T S S K T V L V F L N L I F W G A A G I L C Y V G A Y V F I	
ACTTATGATGACTATGACCACTTCTTTGAAGATGTGTACACGCTCATCCCTGCTGTAGTGATCATAGCTGTAGGAGCCCTGCTTTTCATC	
TGAATACTACTGATACTGGTGAAGAAACTTCTACACATGTGCGAGTAGGGACGACATCACTAGTATCGACATCCTCGGGACGAAAAGTAG	270
T Y D D Y D H F F E D V Y T L I P A V V I I A V G A L L F I	
ATTGGGCTAATTGGCTGCTGTGCCACAATCCGGGAAAGTCGCTGTGGACTTGCCACGTTTGTCATCATCCTGCTCTTGGTTTTTGTCACA	
TAACCCGATTAACCGACGACGACGGTGTTAGGCCCTTTCAGCGACACCTGAACGGTGCAAACAGTAGTAGGACGAGAACCAAAAACAGTGT	360
I G L I G C C A T I R E S R C G L A T F V I I L L V F V T	
GAAGTTGTTGTAGTGGTTTTGGGATATGTTTACAGAGCAAAGGTGGAAAATGAGGTTGATCGCAGCATTCAGAAAGTGTATAAGACCTAC	450
CTTCAACAACATCACCAAAACCCTATACAAATGTCTCGTTTCCACCTTTTACTCCAACTAGCGTCGTAAGTCTTTCACATATTCTGGATG	730
É V V V V L G Y V Y R A K V E N E V D R S ! Q K V Y K T Y	
AATGGAACCAACCCTGATGCTGCTAGCCGGGCTATTGATTATGTACAGAGACAGCTGCATTGTTGTGGAATTCACAACTACTCAGA C TGG	
TTACCTTGGTTGGGACTACGACCGATCGGCCCGATAACTAATACATGTCTCTGTCGACGTAACAACACCTTAAGTGTTGATGAGTCTGACC	540
NG INFUAASKAIUTYUKULHEEGIHNISUW	
GAAAATACAGATTGGTTCAAAGAAACCAAAAACCAGAGTGTCCCTCTTAGCTGCTGCAGAGAGACTGCCAGCAATTGTAATGGCAGCCTG	630
CTTTTATGTCTAACCAAGTTTCTTTGGTTTTTGGTCTCACAGGGAGAATCGACGACGTCTCTCTGACGGTCGTTAACATTACCGTCGGAC	
ENTOWFKETKNQSVPLSCCRETASNCNGSL	
GCCCACCCTTCCGACCTCTATGCTGAGGGGTGTGAGGGCTCTAGTTGTGAAGAAGCTACAAGAAATCATGATGCATGTGATCTGGGCCGCA	720
CGGGTGGGAAGGCTGGAGATACGACTCCCACACTCCGAGATCAACACTTCTTCGATGTTCTTTAGTACTACGTACACTAGACCCGGCGT	120
A H P S D L Y A E G C E A L V V K K L Q E I M M H V I W A A	
CTGGCATTTGCAGCTATTCAGCTGCTGGGCATGCTGTGTGCTTGCATCGTGTTGTGCAGAAGGAGTAGAGATCCTGCTTACGAGCTCTTC	
GACCGTAAACGTCGATAAGTCGACGACCCGTACGACACACGAACGTAGCACAACACGTCTTCCTCATCTCTAGGACGAATGCTCGAGAAG	810

43 HLTAH80	Page 2
ATCACTGGCGGAACCTATGCATAGTTGACAATCTCAAGCCTGAGCTTTTTGGTCTTGTTCTGATTTGGAAGGTGAATTGAGCAGGTCT	
TAGTGACCGCCTTGGATACGTATCAACTGTTAGAGTTCGGACTCGAAAAACCAGAACAAGACTAAACCTTCCACTTAACTCGTCCAGAC	+ 900 :G
ITEGTYA. (SEGIDNO: 23)	
TGCTGTTGGCCTCTGGAGTTCATTTAGTTAAAGCACATGTACACTGGTGTTGGACAGAGCAGCTTGGCTTTTCATGTGCCCACCTACTT	
ACGACAACCGGAGACCTCAAGTAAATCAATTTCGTGTACATGTGACCACAACCTGTCTCGTCGAACCGAAAAGTACACGGGTGGATGAA	→ 990 T
CCTACTACCTGCGACTTTCTTTTTCCTTGTTCTAGCTGACTCTTCATGCCCCTAAGATTTTAAGTACGATGGTGAACGTTCTAATTTCA	.G + 1080
GGATGATGGACGCTGAAAGAAAAAGGAACAAGATCGACTGAGAAGTACGGGGATTCTAAAATTCATGCTACCACTTGCAAGATTAAAGT	
AACCAATTGCGAGTCATGTAGTGTGGTAGAATTAAAGGAGGACACGAGCCTGCTTCTGTTACCTCCAAGTGGTAACAGGACTGATGCCG	A + 1170
TTGGTTAACGCTCAGTACATCACACCATCTTAATTTCCTCCTGTGCTCGGACGAAGACAATGGAGGTTCACCATTGTCCTGACTACGGC	
AATGTCACCAGGTCCTTTCAGTCTTCACAGTGGAGAACTCTTGGCCAAAGGTTTTTGGGGGGGAGGAGGAAACCAGCTTTCTGGTTA	.A + 1260
TTACAGTGGTCCAGGAAAGTCAGAAGTGTCACCTCTTGAGAACCGGTTTCCAAAAACCCCCCTCCTCCTCCTTTGGTCGAAAGACCAAT	
GGTTAACACCAGATGGTGCCCCTCATTGGTGTCCTTTTAAAAAATATTTACTGTAGTCCAATAAGATAGCAGCTGTACAAAATGACTAA	A + 1350
CCAATTGTGGTCTACCACGGGGAGTAACCACAGGAAAATTTTTTATAAATGACATCAGGTTATTCTATCGTCGACATGTTTTACTGATT	
ATAGATIGTAGGATCATATGGCGTATATCTTGGTTCATCTTCAAAATCAGAGACTGAGCTTTGAAACTAGTGGTTTTTAATCAAAGTTG	G + 1440
TATCTAACATCCTAGTATACCGCATATAGAACCAAGTAGAAGTTTTAGTCTCTGACTCGAAACTTTGATCACCAAAAATTAGTTTCAAC	С
TTTATAGGAGGAGTATAATGTATGCACTACTGTTTAAAAGAATTAGTGTGAGTGTGTTTTTGTATGAATGAGCCCATTCATGGTAAG	T + 1530
GAAATATCCTCCTCATATTACATACGTGATGACAAAATTTTCTTAATCACACTCACACAAAAACATACTTACTCGGGTAAGTACCATTC	
TTAAGCTIGTIGGAAATAATGTACCCATGTAGACTAGCAAAATAGTATGTAGATGTGATCTCAGTTGTAAATAGAAAAATCTAATTCA	A + 1620
GAATTCGAACAACCTTTATTACATGGGTACATCTGATCGTTTTATCATACATCTACACTAGAGTCAACATTTATCTTTTTAGATTAAGT	
TAAACTCTGTATCAGCCCCCAACAAAAAAAAAAAAAAAA	
ATTIGAGACATAGTCGGGGGTGTTTTTTTTTTTTTTTTTT	

Angele angele and the second of the second o

11

Page 1

HTPBAZT											
CACGAGCGCAGAGCTTGGGGGCTTCCTTGGTCGCACCCACC	90										
GTGCTCGCGTCTCGAACCCCGAAGGAACCAGCGTGGGTGG	90										
TCTTTCCTGTGGCCAGCCCAGAACTGAAGCGCTGCGGCATGGCGCGCGC											
AGAAAGGACACCGGTCGGGTCTTGACTTCGCGACGCCGTACCGCGCGGGACGGAC	180										
M A R A C L Q A V K Y L M F A F N											
TGTTCTTCTGGCTGGGAGGCTGTGGCGTGCTGGGTGTCGGCATCTGGCTGG											
ACAAGAAGACCGACCCTCCGACACCGCCACAGCCGTAGACCGACC	270										
LFFWLGGCGVGIWLAATOGSFATLSSS											
TCCCGTCCCTGTCGGCTGCCAACCTGCTCATCATCACCGGCGCCTTTGTCATGGCCATCGGCTTCGTGGGCTGCCTGGGTGCCATCAAGG											
AGGGCAGGGACGGCTTGGACGAGTAGTAGTGGCCGCGGAAACAGTACCGGTAGCCGAAGCACCCGACGGACCCACGGTAGTTCC	360										
FPS LS AAN LLIITGAF V MAIGF V GC LGAIK											
AGAACAAGTGCCTCCTGCTCACTTTCTTCCTGCTGCTGCTGCTGGTGTTCCTGCTGGAGGCCACCATCGCCATCCTCTTCTTCGCCTACA	450										
TCTTGTTCACGGAGGACGAGTGAAAGAAGGACGACGACGACCACAAGGACGACCTCCGGTGGTAGCGGTAGGAGAAGAAGCGGATGT											
ENKCLLLTFFLLLLVFLLEATIAILFFAY											
CGGACAAGATTGACAGGTATGCCCAGCAAGACCTGAAGAAAGGCTTGCACCTGTACGGCACGCAGGGCAACGTGGGCCTCACCAACGCCT											
GCCTGTTCTAACTGTCCATACGGGTCGTTCTGGACTTCTTTCCGAACGTGGACATGCCGTGCGTCCCGTTGCACCCGGAGTGGTTGCGGA	540										
T D K I D R Y A Q Q D L K K G L H L Y G T Q G N V G L T N A											
GGAGCATCATCCAGACCGACTTCCGCTGCTGTGGCGTCTCCAACTACACTGACTG											
	630										
WSIIQTDFRCCGVSNYTDWFEVYNATRVPD											
CCTGCTGCTTGGAGTTCAGTGAGAGCTGTGGGCTGCACGCCCCGGCACCTGGTGGAAGGCGCCGTGCTACGAGACGGTGAAGGTGTGGC	720										
GGACGACGACCTCAAGTCACTCTCGACACCCGACGTGCGGGGGCCGTGGACCACCTTCCGCGGCACGATGCTCTGCCACTTCCACACCG											
SCCLEFSESCGLHAPGTWWKAPCYETVKVW											
TTCAGGAGAACCTGCTGGCTGTGGGCATCTTTGGGCTGTGCACGGCGCTGGTGCAGATCCTGGGCCTGACCTTCGCCATGACCATGTACT	L										
AAGTCCTCTTGGACGACCGGACACCCGTAGAAACCCGACACGTGCCGCGACCACGTCTAGGACCCGGACTGGAAGCGGTACTGGTACATGA	810										
LOENLLAVGIFGLCTALVOILGLTFAMTMY											
GCCAAGTGGTCAAGGCAGACACCTACTGCGCGTAGGCCGCCCACCGCCGGCTTCTCTGCCAAAAGGACGCCCACGGGGAGATGGCCGCAC											
CGGTTCACCAGTTCCGTCTGTGGATGACGCGCATCCGGCGGTGGCGGCCGAAGAGACGGTTTTCCTGCGGGTGCCCCTCTACCGGCGTG	900										
_											
COVVKADTYCA. (SEQID'NO: 24)											

Page 1

																							AH	II	\mathcal{I}	15	3			
GCG	CCC	SCCC	GGC	CGC	AGC	ATO	GGG	CGC	TT	CCG					GTGC						CT	TGG	TTC	AAC	CTO	3CTC	TTC	TGG		90
CGC	GGC	GGG	ccc	GCG	TCC	TÀC	ccc	GCG	AA	GGC					CACG						GA	ACC	AAG	TTG	GAC	GAC	SAAG	ACC		30
						м	G	R	F	R	G	G	L	R	С	I	K	Y	L	L	L	G	F	N	L	L	F	w	L	•
GCT	GG/	тсс	GCC	GTC	ATI	GCI	TTT	GGA	ACT.	ATGO	3111				AGGT			AAA	GGA	STTA	TC	ATC	AGAG	GAC	AAC	STCC	CCA	GAG	TAT	180
CGA	CCI	rago	CGG	CAG	TAA	CGA	AAA	CCT	GA	TAC	AAA				TCCA			m	CCT	CAAT	AG	TAGI	стс	CTG	TTC	CAGC	GGT	CTC	ATA	100
Α	G	s	Α	٧	I	A	F	G	L	W	F	R	·F	G	G	A	ī	κ	Ε	L	s	s	Ε	٥	K	S	Р	Ε	Y	
TTC	TAT	GTG	GGG	CTG	TAT	GTT	CTG	GTT	GG	4GC(GGG	GCC	CTG	ATO	SATG	GCC	GTO	GGG	STT	TTC	:GG	ATGO	TGC	GGA	GCC	CATO	CGG	GAG	TCG	
			-i	+											TAC	-+-													+	270
_	v	V			v	v	1	v	c		·						.,	_	_	_	_	_	_	_		u		_		
r	ľ	•	G	_	'	•	L	٧	G	А	G	А	L	n	rı	А	٧	G	r	Г	G	L	L	G	А	n	ĸ	E	5	
CAA	TGT	GTO	CTT	GGA	TCA	TTT	TTT	ACC	TG	CTC	CTG	GTG	ATA	TT	rgct	GCT	GAA	GT	AAC	ACT	GG	AGTA	TTT	GCT	TTI	ATA	GGC	AAG	GGG	360
GTT	ACA	CAC	GAA	CCT	AGT	AAA	AAA	TGG	AC	GGAG	GAC	CAC	TAT	ΆΑΑ	ACGA	CGA	CTI	rca:	TTG	TGA	CC.	TCAT	ΑΑΑ	CGA	AAA	ATĀT	CCG	TTC	ccċ	500
a	С	v	L	G	S	F	F	T	С	L	L	٧	I	F	A	A	Ε	٧	T	T	G	٧	F	A	F	1	G	κ	G	
GTA	GCT	ATC	ÇGA	CAT	GIT	CĄG	ACC	ATG	TA	rga A	GAG	GCT	TAC	AA	TGAT	TAC	CTI	TAA.	AGAC	AGG	GG	444	GGC	AAT	GGG	SAÇA	стс	ATC		"50
CAT	CGA	TAG	GCT	GTA	CAA	GTC	TGG	TAC	ATA	CTI	rc Tc	CGA	ATG	TT	ACTA	ATG	GAA	TT.	гст	тсс	cc.	TTT	CCG	TTA	ccc	TGT	GAG	TAG		450
٧	A	ı	R	Н	٧	a	T	M	Y	Ε	Ε	Α	Y	N	D	Y	L	κ	D	R	G	к	G	N	G	T	L	ī	т	
TTC	CAC	TCA	ACA	TTT	CAG	TGC	TGT	GGA	AAA	GAA	AGC	TCC	GAA	CAC	GTC	CAA	сст	ACA	ATGO	CCA	AAC	GAG	CTT	CTA	GGA	ACAC	AAG	AAT	TGC	
AAG	GTG	AGT	TGT	AAA	GTC	ACG	ACA	CCT	TTI	CTT	TCG				CAG		GGA	TG	race	GGT	TTO	стс	GAA	GAT	CCT	GTG	TTC	TTA	ACG	540
F	н	s	Τ.	F	Q	С	С	G	к	Ε	s	s	E	٥	v	Q	Р	Т	С	Р	K	ε	L	L	G	н	ĸ	N	С	
ATC	GÄT	GAA	ATT	GAG	ACC	ΑΤΔ	ΔΤΟ	ΔGT	GTI	ΔΔΩ	יר דר		CTC	ΔΤΤ	GGA	ΔΤΤ	GTC	.cc.	ΓΔΤΊ	.cca	ΔΤΊ	rcra	GGT	CTG	ΔCG	חדמ	TTT	ດຄວ	ΔTG	
			+			-+-		+					+		CCT	-+-					-+-		+						+	630
	•		177	•					-			316	GAG	11 44		1 44	CAG		1100		100	1001		GAC	100					
1	D	Ε	I	ε	Τ	I	1	S	٧	K	L	0	L	1	G	i	٧	G	I	G	I	Α	G	L	T	1	F	G	M	
ATA	TTC	AGC	ATG	GTC	стс	TGC	TGT	GCG	ATA	CGA	AAC	TCA	CGA	GAT	GTG	ATA	TGA	A GC	TAC	TTC	TAC	ATG	AAA	ATT	GCA	ATC	TAA	AGC	TTŢ	700
TAT	AAG	TCG	TAC	CAG	GAG	ACG	ACA	CGC	TAT	GCT	TTG	AGT	GC T	CTA	CAC	TAT.	ACT	TCG	ATG	AAG	A TG	TAC	TTT	TAA	CGT	TAG	ATT	TCG		720
I	F	s	М	٧	L	С	С	Α	i	R	N	s	R	D	V	1		(5	CC	ΣŔ		7(,O	<u>'</u> 2	5	1				
		AAA	+	->	734	C	Sa	P	I	D	N):	7),	, .											ر				

Päge 1

HAIDO59	
AGTGTTTATGGGACTAAAAAACTTTTAACACCTTTTTAGGGGAAATATTTTGGTCCTATACAAAACATGTAAATATGCTTTATTACTTTC	90
TCACAAATACCCTGATTTTTTGAAAATTGTGGAAAAATCCCCTTTATAAAACCAGGATATGTTTTGTACATTTATACGAAATAATGAAAG	90
ATTTTCTGACCCTGCTGTAAACTACTGCAACCCTCACATCCCTCAAAGGGACTTTTATGTCAAACTCTTCTGTTTCTCCAAATATAAGGA	. 180
TAAAAGACTGGGACGACATTTGATGACGTTGGGAGTGTAGGGAGTTTCCCTGAAAATACAGTTTGAGAAGACAAAGAGGTTTATATTCCT	100
AAAAAGACTAAAGCAAGAGATCTGGCAGTTGAAAATTGTGGGAAAGAGAATTTGTATGGGCACTGTATCTATGAAATACCTCATACTTAC TTTTTCTGATTTCGTTCTCTAGACCGTCAACTTTTAACACCCTTTCTCTTAAACATACCCGTGACATAGATACTTTATGGAGTATGAATG	270
TTTTTCTGATTTCGTTCTCTAGACCGTCAACTTTTAACACCCTTTCTCTTAAACATACCCGTGACATAGATACTTTATGGAGTATGAATG	2,0
GTTTACATGTTTTCCTAACTTTTTGTATTTTTCTTGTATAGCCACCTAGAGAATTCTTCATAGATTAAGAACTACAGTTTTCACCACTTA	360
CAAATGTACAAAAGGATTGAAAAAACATAAAAAGAACATATCGGTGGATCTCTTAAGAAGTATCTAATTCTTGATGTCAAAAGTGGTGAAT	500
ACATAAGTAAAACAAAGTCCTTCATAATTTAACCATTAGCATCTTTGGCCAAACCAAAATAAAGAAAAGCATCTTCTCCTAGTTGTGTGT TGTATTCATTTTGTTTCAGGAAGTATTAAATTGGTAATCGTAGAAACCGGTTTGGTTTTATTTCTTTTCGTAGAAGAGAGGATCAACACACA	450
TGTATTCATTTTGTTTCAGGAAGTATTAAATTGGTAATCGTAGAAACCGGTTTGGTTTTATTTCTTTTCGTAGAAGAGGATCAACACACA	430
GGGCAACAGAAACAAGTTAAGGAAACAAAAATACTTATATATA	540
CCCGTTGTCTTTGTTCAATTCCTTTGTTTTTATGAATATATGTGTCTTGTTTTTATTACAAGAAAAATACGTTTAGGGGACACTTTTA	340
AAAATTITCAATGITTAAAAAAAAAAAAAAAAAAAAAAAA	
AAAATTITCAATGITTAAAAAAAAAAAAAAAAAAAAAAAA	

Same Age :

																<u> </u>												4		
			+-										-}													-+-			CGG	90
AAG	CCG	TGC	TC	SACO	SC C C	GCC	CACC	CGC	CGA	ACCO	CGCC	CGGG	GCC	CTO	GGC	GCC	AGA	GTC	AGA	AGAG	ATC	CGC	GTO	AGG	GAA	GCG	GCG	AAG	GCĊ	
																														•
AGC	ccc	TGG	CAC	GGC	CCA	GAA	GCC	ATG	GCC	CAC	TAI	TAAG	ACT	GAC	CAC	GAC	GAC	TGG	CTO	ATC	ATC	TAC	TTG	AAG	TAT	TTA	CTC	:	GTC	180
TCG	GGG	ACC	GTO	cce	GGT	CTI	rcgo	STAC	CGC	GTC	SATA	47 T C	TGA	CTO	CGTC	CTO	CTG	SACC	GAC	TAG	TAG	ATG	AAC	TTC	ATA	AA1	GAG	AAA	CAG	
								M	A	Н	Y	K	τ	Ε	Q	а	а	W	Ĺ	i	i	Y	L	κ	Y	L	Ł	F	V	
TTC	AAC	TTC	TTC	TTC	TGG	GŢC	GGG	GGA	GC/	GCC	GTC	CTG	GCT	GTO	GGG	CAŢC	TGG	ACC	сто	GTG	GAG	AAG	ĄGT	GGC	TAC	CTC	AGC	GTC	CTG	
AAG	TTG	AAG	AAG	AAG	ACC	CAC	CCC	CCT	CGI	CGG	CAG	GAC	CGA	CAC	ccc	STAC	ACC	TGG	GAC	CAC	стс	TTC	TCA	CCG	ATG	GAG	TCG	CAG	GAC	270
. F	N	F	F	F	W	ν	G	G	Α	Α	٧	L	A	v	G	i	W	т	L	٧	ε	ĸ	s	G	Y	L	s	v	L	
GCC	TCC	AGC	ACC	TTT	GCC	GCC	TCC	GCC	TAC	ATC	сто	ATC	TTI	GCG	GGC	GTA	стт	GTC	ATG	GTG	ACC	GGC	TTC	CTG	GGC	ттс	GGT	GCC	ATC	
CGG.	AGG	TCG	TGG	AAA	CGG	CGG	AGG	CGG	ATO	TAG	GAG	TAG	AAA	CGC	CCC	CAT	GAA	CAG	TAC	CAC	TGG	CCG	AAG	GAC	CCG	AAG	CCA	CGG	TAG	360
Δ	s	S	т	F		А	s	Δ	Y	Ţ	i	ī	F	Δ	G	v	1	v	м	v	т	G	F	1	G -	E	c	Δ	1	
.,	Ĭ	Ů	•	•			Ū	.,	·	٠	_	•	,		J	·	_	•	• • •	•	•	J	•	_	٠		٠		•	
			+	+				+					+-					+		GTT			+			-,-			+	450
GAG	4CC	CTC	GCC	TTC	CCG	ACG	GAG	AGG	TGC	ATA	AAG	ACG	GAC	CAAC	GAG	CAG	TAG	AAG	GAC	CAA	СТС	GAC	CAC	CGC	CCT	CAG	GAC	CGG	GTA	
L	W	ε	R	K	G	С	L	s	T	Y	F	С	L	L	L	٧	I	F	L	٧	E	L	٧	A	G	٧	. L	A	Н	
GTG	ГАТ	TAC	ÇAG	AGG	CTG	AGT	GAT	GAA	ст	AAG	CAG	CAC	TTG	AAC	CGG	ACT	CTG	GCT	GAG	AAC	TAC	GGG	ÇAG	CCG	GAG	CAC	GCA	GAT	CAÇ	
CAC	λTA	ATG	GTC	TCC	GAC	TCA	CTA	СТТ	GAC	TTC	GTC	GTG	AAC	TTG	GCC	TGA	GAC	CGA	CTC	TTG	ATG	ccc	GTC	GGC	стс	GTG	CGT	CTA	GTG	540
٧	Y	Y	0	R	L	s	D	Ε	L	K	Q	н	Ł	N	R	T	L	А	ε	N	Υ	G	Q	ρ.	E	н	А	D	н	
GCC.	rca	GTG	GAC	CGA	СТС	CAG	CAG	GAT	TTC	AAG	TGC	TGC	GG A	AGC	AAC	AGC	TCA	GCC	GAC	TGG	CAG	CAC	AGC	ACG	TAC	ATC	CTG	TTG	CGG	
CGG			+			-+-		+	+				+					+	+				 						+	630
Δ	S	v	n	R	,	n	n	n	E	ĸ	'n	٥	c		M	e	c	^		1.7	0	ы	e	т	···· •	1	1	,	p	
	•	•	Ü	.,	_	4	•	Ü		K	٠	C	G	3	13	J	J	^	U	~	u	"	3	•		•	_	₩.		
GAG	-		}			-1-								+														+	+	720
CTC	GG	CTC	C CG	GCG	GTC	CAC	GGG	CTG	TCG	ACG	ACG	TTC	TGT	CAC	CAC	C GC	GCG	ACG	CCG	GTC	GCC	CGG	GTG	GGG	AGG	TTG	T AG	ATA	TTC	
Ε	A	E	G	R	Q	٧	ρ	D	s	С	С	K	T	٧	٧	Α	R	С	G	Q	R	A	н	Р	s	N	ľ	Y	Κ	
GTG	AG	GGA	GGC	TGC	CTC.	ACC	AAG	c TG	GAG	CAG	TTC	CTG	GCC	GAC	CAC	CTG	CTG	CTT.	ATG	GGG	GCA	GTG	GGC	ATC	GGG	GTG	GCC	TGC	CTG	
CACC			+			 -							+					+	 +							-+-		+		810
¥	ε	G	G	С				L					А	D			L	•	м		Δ	٧	. 47	ı	G	٧	Α	С	L	
		~~-															_			-	.,									
CAGA	+			+					+									+	+							-:-			+	900
GTC1	AG.	AUG	LUC	ΙĄĊ	LAA	GAG	r GG.	ACG.	ACG	AAC																		_	LGG	
ũ	i	С	G	M	٧	L	T	С	С	L	Н	Q	R	L	0	R	н	F	Y	. (S	? (Q)	7	D	Ν	0	.2	6		

Page 2 HHFEKHO

CCTCAAGACAACATGTGGCACATGCCATCTGCAAGG
GGAGTTCTGTTGTACACCGTGTACGGTAGACGTTCC

936 (SCQ IDNO: 11)

Page 1

	HGBGV89																													
AGC	TTA	CTT	TCA	CTO	ACC	GÇC	TGT	CCT	TC	TGA	CAC	CTC	ACC	ATO	TGT	ACC	GGA	AAA	TGT	GCC	CGC	TGT	GTG	GGG	ст	TÇC	:CTC	ATT	ACC	90
TCG	AAT	GAA	AGT	GAO	STGG	ccic	ACA	GGA	AGC	SACT	GTG	GAG	TGG	TAC	ACA	TGC	CCI	TTT	ACA	CGG	GCG	ACA	CAC	ccc	GAG	AGC	GAG	TAA	TGG	90
														м	c	т	G	ĸ	ſ	Δ	R	۲	v	c	,	s	ı	1	τ.	
															_	·	Ī		Ū	••	.,	·	·	•	_	·	_	•	·	
CTC	TGC	CTC	GTC	TG	ATT	GTG	GCC	AAC	GCC	CTC	CTG	CTG	GTA	CCT	AAT	GGG	GAG	ACC	TCC	TGG	ACC	AAC	ACC	AAC	CA	CTC	AGC	TTG		180
GAG	ACG	GAG	CAG	ACC	STAA	CAC	CGG	TTG	CGG	GAG	GAC	GAC	CAT	GGA	TTA	ccc	CTC	TGG	AGG	ACC	TGG	TTG	TGG	TTG	GT/	1GAG	TCG	AAC	GTT	
L	С	L	٧	С	t _e	٧	A	N	A	L	L	L	٧	Ρ	N	G	Ε	T	S	W	T	N	٢	N	Н	L	s	L	۵	
GTC	TGG	сто	ATG	GGC	GGC	TTC	ATT	GGC	GGC	GGC	CTA	ATG	GTA	ста	TGT	CCA	GGG	ATT	GCA	GCC	GTT	CGG	GCA	GGG	GGC	CAAG	GGC	TGC	TGT	
CAG	ACC	GAG	TAC	CCC	ccc	AAG	TAA	CCG	CCC	CCG	GAT	TAC	CAT	GAC	ACA	GGT	CCC	TAA	CGT	CGG	CAA	GCC	CGT	ccc	cc	TTC	CCG	ACG	AC A	270
٧	W	Ł	М	G	G	F	ı	G	G	G	L	M	٧	L	С	Ρ	G	i	A	A	٧	R	A	G	G	ĸ	G	С	С	
COT	ec1	ecc.	TGC	TCI	ree A	Δ Δ.Γ	LEL	TCC	ACC	: A T C	er te	ירפר	TCC	e TC	TTC	TCC	TCC	:c.c.	TTC	ccc	CTC	CTT	CCT	ecc	ΔΤΩ	- T A C	דכ			
	<u> </u>		+			+-							+		AAG	-							+							
CCA	CGA		ACG	MLA	(,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	110	GCG	ALC			JAU	.60.6	AGC	CAU	AAG	AGG	AGC		AAG		LAL	GAA	CCA	CGG	IAC	1A 1 G	исс	e		
G	A	G	С	С	G	N	Ŕ	С	R	M	L	R	S	ν	F	S	S	A	F	G	٧	L	G	A	I	Υ	С	L		
GTG	TC T	GGA	GCT	GGG	сто	CGA	AAT	GGA	ccc	AGA	TGC	TTA	ATG	SAAC	GGC	GAG	TGG	GGC	TAC	CAC	TTC	GAA	GAC	ACC	GCC	GGA	GCT	TAC		
CAC	AGA	CCT	CGA	ccc	GAG	GCT	TTA	CCT	GGG	TCT	ACG	AAT	TAC	TTG	CCG	CTC	ACC	CCG	ATG	GTG	AAG	CTT	CTG	TGG	CGC	CCT	CGA	ATG		450
٧	s	G	A	G	L	R	N	G	Р	R	С	L	м	N	G	Ε	W	G	Y	н	F	Ε	0	т	Α	G	Α	Y	L	
																_														
			+			-+-		+	+				+		GTC	-+-		+	+				+						-+	540
GAG	TTG	GCG	TGA	GAT	ACC	CTA	GCC	ACG	CTC	CGC	GGG	GGA	GCG	CAC	CAG	GGG	ACC	TTA	CAC	TGC	GAG	AAG	AGC	GAC	GAC	CAC	CGG	CGG.	AGG	
L	N	R	T	L	W	D	R	С	Ε	A	Ρ	P	R	٧	٧	Ρ	W	N	٧	T	L	F	s	L	L	٧	A	Α	s	
TGC	сто	GAG	ATĀ	GTA	CTG	TGT	GGG	ATC	CAG	CTG	GTG	AAC	GCG	ACC	ATT	GGT	GTC	TTC	TGC	GGC	GAT	TGC	AGG	AAA	AAA	CAG	GAC	ACA	сст	
			+			-+-		+					+		TAA								+							630
		_																								_	_	_	_	
С	L	E	I	٧	L	С	G	ì	Ω	Ĺ	٧	N	A	Т	ī	G	٧	F	С	G	D	С	R	K	K	0	υ	1	P	
CAC	TGA	GGC	ŢCC	ACT	GAC	cġc	CGG	GTT	ACA	CCT	GCT	ССТ	ŢCC	TGG	ACG	сŢС	AC T	ccc	TTG	CTC	GCT	AGA	ΑTΑ	AAC	TGC	TTT	GCG	CTC	TCA	720
GTG	ACT	CCG	AGG	TGA	CTG	GCG	GCC	CAA	TGT	GGA	CGA	GGA	AGG	ACC	TGC	GAG	TGA	GGG	AAC	GAG	CGA	TCT	TAT	TTG	ACG	AAA	CGC	GAG		120
н	٠ (5	eG	J	D	N	0:	2	7)																				
AAA					AAC				_		<u>ب</u> ب	<u> </u>	شرى	١.	\sim	7														
		TTT	+		>	73	8	ω	Œ	، لا	レ し	ノい	٧C	<i>'</i> :	12	J														

Page 1

H UVB380	
GGCACGAGAGATTGTCGGCTGCGGGTATATTCCAATTCCCCGTCTCCTCATGAATATGAAGTGAAGGGCTCTGACCCTGGAAGTGGTTC	T → 90
CCGTGCTCTCTAACAGCCGACGCCCATATAAGGTTAAGGGGCAGAGGAGTACTTATACTTCACTTCCCGAGACTGGGACCTTCACCAAG	
AAGCAGGGCAAAATGGGGTCTCGGAAGTGTGGAGGCTGCCTAAGTTGTTTGCTGATTCCGCTTGCACTTTGGAGTATAATCGTGAACAT	
TTCGTCCCGTTTTACCCCAGAGCCTTCACACCTCCGACGGATTCAACAACGACTAAGGCGAACGTGAAACCTCATATTAGCACTTGTA	-+ 180 \T
M G S R K C G G C L S C L I P Ł A L W S I I V N I	
TTATTGTATTTCCCGAATGGGCAAACTTCCTATGCATCCAGCAATAAACTCACCAACTACGTGTGGTATTTTGAAGGAATCTGTTTCTC	:A + 270
AATAACATAAAGGGCTTACCCGTTTGAAGGATACGTAGGTCGTTATTTGAGTGGTTGATGCACACCATAAAACTTCCTTAGACAAAGAC	
LLYFPNGOTSYASSNKLTNYVWYFEGICFS	;
GGCATCATGATGCTTATAGTAACAACAGTTCTTCTGGTACTGGAGAATAATAACAACTATAAATGTTGCCAGAGTGAAAACTGCAGCAA	\A
CCGTAGTACTACGAATATCATTGTTGTCAAGAAGACCATGACCTCTTATTATTGTTGATATTTACAACGGTCTCACTTTTGACGTCGTT	-+ 360 T
G I M M L I V T T V Ł L V L E N N N N Y K C C Q S E N C S K	
AAATATGTGACACTGCTGTCAATTATCTTTCTTCCCTCGGAATTGCTTTTTCTGGATACTGCCTGGTCATCTCTGCCTTGGGTCTTGT	°C + 450
TTTATACACTGTGACGACAGTTAATAGAAAAGAAGGGGAGCCTTAACGAAAAAGACCTATGACGGACCAGTAGAGACCCAGAACA	
K Y V T L L S I I F S S L G I A F S G Y C L V I S A L G L V	!
CAAGGGCCATATTGCCGCACCCTTGATGGCTGGGAGTATGCTTTTGAAGGCACTGCTGGACGTTTCCTTACAGATTCTAGCATATGGAT	T 5110
GTTCCCGGTATAACGGCGTGGGAACTACCGACCCTCATACGAAAACTTCCGTGACGACCTGCAAAGGAATGTCTAAGATCGTATACCTA	.A
Q G P Y C R T L D G W E Y A F E G T A G R F L T D S S I W I	
CAGTGCCTGGAACCTGCACATGTTGTGGAGTGGAACATCATTTTATTTTCCATTCTCATAACCCTCAGTGGGCTTCAAGTGATCATCTG	
GTCACGGACCTTGGACGTGTACAACACCTCACCTTGTAGTAAAATAAAAGGTAAGAGTATTGGGAGTCACCCGAAGTTCACTAGTAGAC	+ 630 :G
Q C L E P A H V V E W N I I L F S I L I T L S G L Q V I I C	:
CTCATCAGAGTAGTCATGCAACTATCCAAGATACTGTGTGGAAGCTATTCAGTGATCTTCCAGCCTGGAATCATTTGAATAAGGACAAA	Α.
	+ 720
LIRVVM Q L S K I L C G S Y S V I F Q P G I I . (SCO) ID	NO:28
TGTTTTCCATTATCAAGACATGGCCATCTATCTAAATATTATATCAACTGTGTTAGACTTGAGGGCAATATTGAAAAATGATGGTGCTTT	
ACAAAAGGTAATAGTTCTGTACCGGTAGATAGATTTATAATATAGTTGACACAATCTGAACTCCCGTTATAACTTTTACTACCACGAAA	+ 810 G
TGCATTIGGTGTTTATTIGTAAAAAATTIGCAGTCCTCACTGCACATGCAAGTATACCACCCTTCCATTTAGTATGTTTTTTAAGTAAT	+ 900
ACGTAAACCACAAATAAACATTTTTTTAAACGTCAGGAGTGACGTGTACGTTCATATGGTGGGAAGGTAAATCATACAAAAAATTCATTA	T

Paige 2

AGAAACCTITATGCAATATGTATATTGCAACATTATTTAATATTCTGGAAAATTGGAAACACCCCAAAATTCTAACCTCAAA

TCTTTGGAAATACGTTATACATATAACGTTGTAATAAAATTATAAGACCTTTTAACCTTTTGGGGGTTTTAAGATTGAGATTGAGTTT

Page 1

HJACE54

TTTG	TG	GAG	GGC	AGC	AGA	GAG	TAI	CCCA	AGCT	GG	ACA	TCCI	TTC	CTG	CTG	ATG	AGC	CCC	AGG	CTG	GAG	GTG	CCC	TGC	TCA	CAT	GCT	CTT		90
AAAC	AC	CTC	CCG	TCG	TCT	CTC	AT	GGGT	CGA	CC.	TGT	AGGA	AAG	GAC	GAC	TAC	TCG	GGG	TCC	GAC	CTC	CAC	GGG	ACG	AGT	GTA	CGA	GAA		•
																м	s	Р	R	Ļ	Ε	٧	Р	С	s	H	A	L	Р.	
																				_									-	
CAGG	<u>.</u>		+			-+-							-+					+	+				+						+	180
GTCC	CA	GAG	AGC	GGA	CCC	GTC	CAC	GTAG	STAT	CA	rgcı	CCCT	GAC	CAG	AAC	GTT	стс	GGC.	TTC	GŢA	AAA.	TGA	CAC	TCG	GAC	TCC	CTG	GTC	CGA	
O	G	L	s	P	G	۵	٧	1	i	٧	R	G	L	٧	L	۵	Ε	Ρ	K	Н	F	T	٧	S	L	R	D	۵	A	
SCCC	ΑT	GC T	CCT	STG	ACA	CTC	AGO	SGCC	TCC	TTO	CGC	AGAC	AGA	ACT	CTG	GCC	TGG	ATC:	TCC	CGC	TGG	GGG	CAG	AAG	AAA	CTG	ATC	TCA	GCC	
CGGG			+	+		- ;-			+				+			-+-		+	+				+						+	270
					_		_			_		_	_	_					•	_		_			.,					
Α	H	Α	Р	٧	Ţ	L	R	Α	S	,	Α	D	R	1	L	А	W	1	5	ĸ	W	G	u	K	X	Ĺ	ı	\$	А	
CCT	TC	стс	TTT	TAC	ccc	CAG	AG/	ATTO	TIT	GAC	GTO	CTG	CTC	CTG	TTC	CAG	GAG	GGA	GGG	CTG	AAG	CTG	GCG	CTC	AAT	GGG	CAG	GGG	CTG	360
GGA	AG	GAG	AAA	A T G	GGG	GTC	TC1	ΓΑΑΘ	AAA	CTO	CAC	GAC	GAG	GAC	AAG	GTC	CTC	CCT	ccc	GAC	TTC	GAC	CGC	GAG	TTA	CCC	GTC	CCC	GAC	
Р	F	L	F	Υ	P	a	R	F	F	ε	٧	L	L	L	F	Q	Ε	G	G	Ļ	κ	L	A	L	N	G	۵	G	Ĺ	
SGGG		۸۲۲	۸۵۲		A A C	C 4 C	CAC	יר רר	r Te	CAC	. r . r	יר דר	rrc	CAC	רדר	ccc	ATC	ACT	CCA	ACT	CTC	CVC	r Tr	TAC	TOT	CTC	רער	TCC.	TCA	
CCC	+		+			-+-			+				+			-+-		+	-+				+						-+	450
	44	100	100	IAC					idac			.0.			0.70	acc	170				CAG		00			0.0	0.0		SCG	OI
G	A	T	S	M	N	a	a	A	Ł	Ε	a	L	R	Ε	L	R	ı	S	G	S	٧	a	L	Υ	С	٧	н	5	•	NO:-2
AGGA	TG	GTT	CCA	GGA	AAT	ACC	GCA	GAA	AAC	AAG	AGI	CAG	CCA	стс	ССС	AGG	GC C	CCA	CTC	TCC	TCC	сст	CAT	TAA	ACC	ATC	CAC	CTG	AAC	540
CCT	AC	CAA	GGT	CT	TTA	TGG	CGT	CTI	TTG	TTC	TCA	GTC	GGT	GAG	GGG	TCC	CGG	GGT	GAG	AGG	AGG	GGA	GTA	ATT	TGG	TAG	GTG	GAC	TTĠ	340
																									•					
CCA	GC.	ACA	TCA	GG	сст							CGA							TTT	GGG	CCT	GAG	GGA	AGG	CAC	AAG	AGT	GCA/	AAG	630
GGT	CG	TGT	AGT	cc	GGA														٩AA	ССС	GGA	CTC	сст	TCC	GTG	TTC	TCA	CGT		
																	_													
STTC	CTI	CGA	ACTO	TG	CAC	CTT						TGG						GCCT	TTC.	AGG	GCC	TGG	ACT	GCA	стс	ACA	GAG	GCA	AGT	720
CAAG	GA	GCT	TGA	SAC	GTG	GAA	GGA	GGT	GGT	ССТ	CGG	ACC	CTA	TAC	CGA	GG T	AGA	CGGA	AAG	TCC	CGG	ACC	TGA	CGT	GAG	TGT	CTC	CGT	TCA	
TTG	+					-+-							+			-+-			-				+			-+-			-	810
CAAC	ATI	CTG	ATTO	TT	TCT	ATG	AGG	TTT	TAT	GTT	ACC	GAA	TTT	CTT	ACA	CCA	GTA	AATA	AAG.	AAA	TAA	TAA	ATA	AAT.	AAA	CAC	CAG	TTT	ATT	
																											_			
TAA	A T	AAG	GTT/	TT	TAT	TTA	AAA	AAA	AAA	AAA	AAA	AAA	AAA.	AAA	AAA	AAA.	AAA	₹ - 86	35	15	26	Ĭ	Ô	NC) :	14	.)			
TATT	TA	TTC	CAAT	AA	ATA	AAT	TTT	TTT	TTT	TTT	TTT	TTT	TTT	TTT	TTT	TTT	TTT	Г				_	-							

HROAD63

Päge 1

CG	GC	тст	TGC	TGT	AGT	CTC	TAC	TCC	TGT	CGT	AAC	GAC	GAG	GAA	CGT	CGG	GAC	CGA	CAC	CGA	TGT	ccc	GGT	CGG	GAA	TGG	GAC	GCG		90
							M	R	T	A	L	L	L	L	A	A	L	A	٧	A	T	Ġ	Ρ	A	L	1	L	R	С	•
:A(GT	GTG	ACC	AGC	тсс	AGC	AAC	TGC	AAG	CAT	тст	GTG	GTC	TGC	CCG	GÇC	AGC	тст	cgç	TTC	TGC	AAG	ACC	ACG	AAC	AÇA	GTG	GAG	CCŢ	1.8
STO	CA	CAC	TGC	TCG	AGG	TCG	TTG	ACG	TTC	GTA	AĞA	CAC	ĊAG	ACG	GGC	CGG	TCG	AGA	GCG	AAG	ACG	TTC	TGG	TGC	TTG	TGT	CAC	СТС	GGA	10
н	. v	С	т	s	s	s	N	С	K	н	s	ν	٧	С	P	A	s	s	R	F	С	κ	T	Т	N	т	γ	Ε	P	
				CCG									t													_ ,				27
GA(TC	CCGA	AGG	GGC	TTT	CAG	ACC	CTG	GTC	CAG	GTC	CAC	CCG	TAC	CTT	ACG	SACT	ACT	GAA	CCT	CGT	CCG	GGG	TGT	CTG	GGG	TGT	СТС	CTA	
L	R	A	s	P	K	V	W	D	a	٧	Q	٧	G	M	Ε	С		(5	Sec	⊋.	II	1 (4C);	3C	$\langle \cdot \rangle$				
SAA	GCI	CAC	ÇCA	CAG	AGG	AŢG	CAG	ccc	CCĄ	GCT	GC A	TGG	ĄAG	GTG	GAG	GĄC	CAGA	AGC	CCŢ	GTG	GAT	CCC	ĊGG	ATT	TCA	CAC	TCC	TTC		
: T	CG	GTGG	GGT	GTC	TCC	TAC	GTC	GGG	GGT	CGA	CGT	ACC.	TTC	CAC	стс	CTG	TCT	TCG	GGA	CAC	CTA	GGG	GCC	TAA	AGT	GTG	AGG	AAG		36
																											,		~=	-7 -1
				T . T		TOT				TOT		TCC		T A A	ATC	A T T	TAA	400									- ($ >$ ϵ	コマ	. 1. 1

HMW6546

Page 1

AGC	GG	300	GA	CC	CTC	GTG.	TGA/	AGGG	STG	CAG	TACC	TAA	GCC	GGA	GCG	GGG	TAG	AGG	CGG	GCC	GGC	ACC	ccc	TTC	TGA	CCT	CCA	GTG		00
TC	CC	GGG	SCT.	rGG	GAG	CAC	ACT.	rcc	CAC	STC	TGG	ATT	CGC	CCI	CGC	ccc	ATC	TCC	GCC	CGG	CCC	TGC	GGG	AAG	ACT	GGA	GGT	CAC		90
	<u> </u>		-					-+					 -										+					CTG		180
GGC	CG	SAG	rtci	'AG	TCTO	STAC	CCGC	GTC	:110	SAAC	TTC	CTG	AAC	CGC	ССТ	GCC	GAC	GGG	CGG	ccc	GGG	GCC	CCG	TAC	CCG	TGC	CGG	GAC	TTC	
						M	A	۵	N	L	K	O	L	A	G	R	L	Ρ	A	G	P	R	G	M	G	T	A	L	κ	
СТС	TTO	C T C	GGG	GC	CGGC	GCC	GTO	GCC	TAC	GGT	GTG	CGC	GAA	TCT	GTG	ттс	ACC	GTG	GAA	GGC	GGG	CAC	AGA	GCC	ATC	TTC	ттс	AATO		
GAC	AAC	GAG	ccc	CG	GCC	CGC	CAC	CGG	ATO	SC C A	CAC	GCG	CTT	AGA	CAC	AAG	TGG	CAC	CTT	CCG	ccc	GTG	TCT	CGG	TAG	AAG	AAG	TTAC		270
L	L	L	G	A	G	A	٧	A	Y	G	٧	R	Ε	s	٧	F	Ţ	٧	Ε	G	G	н	R	A	I,	F	F	N	R	
ATC	GGT	GGA	ĢTE	CA	GCAG	GĄC	ACT	ATC	СТ	GCC	GAG	GGC	стт	CAC	TTC	AGG	ATC	ССТ	TGĢ	TTC	CAG	TAC	ccc	ATT	ATC	TAT	GAC	ATT		200
TAG	CCA	CCI	CAC	GT	GTC	CTO	TGA	TAG	GAC	CGG	CTC	CCG	GAA	GTG	AAG	TCC	TAG	GGA	ACC	AAG	GTC	ATG	GGG	TAA	TAG	ATA	CTG	TAAG		360
ı	G	G	y	a	a	0	T	ı	L	A	Ε	G	L	H	F	R	ı	P	W	F	a	Y	Ρ	ı	-1	Y	0	i	R	
GCC	AG/	CCT	CGA	AA	ATC	TCC	TCC	CCT	ACA	GGC	TCC	AAA	GAC	CTA	CAG	ATG	GTG	AAT.	ATC	тсс	CTG	CGA	GTG	TTG	тст	CGA	ccc	AATG		45
CGG	TCT	GGA	GCT	TT	TTAG	AGG	AGG	GGA	TGI	CCG	AGG	TTT	CTG	GAT	GTC	TAC	CAC	TTA	TAG	AGG	GAC	GCT	CAC	AAC	AGA	GCT	GGG	TTAC	GA	
A	R	P	R	K	I	S	S	P	7	G	s	κ	0	L	a	M	V	N	1	S	L	R	٧	L	s	R	Ρ	N	A	
CAG	GAG	стт	ÇCT	AGC	ATG	TĄC	CAG	CGC	CTA	GGG	CTG	GAC	TAC	GAG	GAA	CGA	GTG	TTG	CCG	тсс	ATT	GTC	AAC	GAG	GTG	CTC	AAG	AGTG	ΣTG	54
GTC	стс	GAA	GGA	TCC	TAC	ATG	GTC	GCG	GAT	ccc	GAC	CTG	ATG	СТС	CTT	GCT	CAC	AAC	GGC	AGG	TAA	CAG	TTG	CTC	CAC	GAG	TTC.	TCAC		341
۵	Ε	L	ρ	s	М	Y	Q	R	L	G	L	0	Y	Ε	Ε	R	V	L	P	s	i	٧	N	ε	٧	L	ĸ	s	y	
GTG	GCC	AAG	TTC	AA T	GCC	TÇA	CAG	CTG	ATC	ACC	ÇAG	CGG	GCC	CAG	GTA	TÇC	CTG	TTG	ATÇ	CGC	CGG	GAG	ÇTG	ACA	GAG	AĢG	GCCA	AAGG	AÇ	~ ~
CAC	cee	TTC	AAG	TTA	CGG	AGT	GTC	GAC	TAG	TGG	GTC	GCC	CGG	GTC	CAT	AGG	GAC	AAC	TAG	GCG	GCC	ctc	GAC	TGT	СТС	TCC	CGG.	TTCC		630
v	A	κ	F	N	A	s	Q	L	i	T	a	R	A	Q	V	s	Ł	L	i	R	R	Ε.	L	T	Ε	R	A	ĸ	Đ	
TTC.	AGC	стс	ĄTC	CTG	GAT	GĄT	GTG	GCC	ATÇ	AÇA	GAG	C TG.	AGC	TTT	AGC	GAC	GAG	TAC	ACAI	GCT	GCT	GTA	ĢAAI	GCC	AAA	CAAI	STG	SCCC	AĢ	700
AAG	TCG	GAG	TAG	GAC	CTA	CTA	CAC	CGG	TAG	TGT	CTC	GAC	TCG	AAA	TCG	GC T	стс.	ATG	TGT	CGA	CGA	CAT	CTT	CGG	TTT	GTT	CACC	GGG		720
F	s	L	1	L	D,	D	٧	A	ı	Т	Ε	L	s	F	s	R	Ε	Y	T	A	A	٧	Ε	Α.	ĸ	۵	٧	A	۵	
CAG	GAG	GCC	ÇAG	CGG	GCC	CAA	TTC	TTG	GTĄ	GAA	AAA	GCA	AAG	CAG	GAA	CAGO	GG	CAG	444	ATT	STG	CAG	GCCI	GAG	GGT	GAG	3C C (SAGG	СŢ	
GTC	CTC	CGG	GTC	GCC	CGG	GTT.	AAG	AAC	CAT	CTT	TTT	CGT	TTC	GTC	CTT	STC	GCC	GTC	ITT.	TAA	CAC	GTC	CGG	CTC	CCA	CTC	GGC	TCC	- :	810
Q	Ε	Α	0	R	A	Q	F	L	٧.	ε	ĸ	A	ĸ	٥	Ε	Q	R	Q	K	ı	٧	۵	A	ε	G	Ε	A	ε	A	
GC C	AAG	ATG	CTT	GGA	GAA	GÇA	CTG.	AGC.	AAG	AAC	CCT	GGC.	TAC.	ATC.	AAAC	:TTC	GC	AAG/	ATŢ	CGA	GCAI	GCC	CAG	AATA	ATC:	TÇC/	4AGA	ACGA	TÇ	900
CGG.	TTC	TAC	GAA	CCT	стт	CGT	GAC	TCG	TTC	TTG	GGA	CCG	ATG.	TAG	TTT	SAAG	GCG.	TTC	ΓΑΑ	GCT	GT	CGG	GTC	TTA	TAG	AGGI	гтст	rgcT		300
																												_		

HMWF546

GCCACATCACAGAATCGTATCTCACAGCTGACAACCTTGTGCTGAACCTACAGGATGAAAGTTTCACCAGGGGAAGTGACAGCCTC
CGGTGTAGTGTCTTAGCATAGATAGAGTGTCGACTGTTGGAACACGACTTGGATGTCCTACTTTCAAAGTGGTCCCCTTCACTGTCGGAG

ATCAAGGGTAAGAAATGAGCCTAGTCACCAAGAACTCCACCCCCACAAGAAGTGGATCTGCTTCTCCAGTTTTTGA

IXGXX. (SOQIDNO: 31)

57 hMF. WOL

Page 1

GGCACGAGATGACATCACTAAGTGGCCGATCTGCACAGAGCAGGCCAGGAGCAACCACACAGGCTTCCTGCACATGGACTGCAGATCAA CCGTGCTCTACTGTAGTGATTCACCGGCTAGACGTGTCTCGTCCGGTCCTCGTTGGTGTCCGAAGGACGTGTACCTGACGCTCTAGTT MDCEIK GGGCCGCCCTGCTGCATCGGCACCAAGGGCAGCTGTGAGATCACCACCCGGGAATACTGTGAGTTCATGCACGGCTATTTCCATGAGGA CCCGGCGGGGACGACGTAGCCGTGGTTCCCGTCGACACTCTAGTGGTGGGCCCTTATGACACTCAAGTACGTGCCGATAAAGGTACTCCT G R P C C I G T K G S C E I T T R E Y C E F M H G Y F H E E AGCAACACTCTGCTCCCAGGTGAGGCGAGGCAGGCCTGGAGTAGTGGAGGAGGGACGCTGGGCATGGCAGCCTGCTGGGGCCGGGGCTC TCGTTGTGAGACGAGGGTCCACTCCGCTCCGGCCCCGACCTCATCACCTCCTCTCCTGCGACCCGTACCGTCGGACGACCCCGGCCCCGAG ATLC SOVRRGRPG V V E E R T L G M A A C W G R G S ACGCACTCCCTCCCATGTCGGAGCCTCAGACTCAGGCTGCTTCTGGGGCGCTGAGCACCATATGCCCATTCCCAGGTGCACTGTTTTGGA TGCGTGAGGGAGGGTACAGCCTCGGAGTCTGAGTCCGACGAAGACCCCGCGACTCGTGGTATACGGGTAAGGGTCCACGTGACAAAACCT R.T.P.S.H.V.G.A.S.O.S.G.C.F.W.G.A.E.H.H.M.P.I.P.R.C.T.V.L.D CAAGGTGTGTTGGGCTGCCTTCCTCAACCCTGAGGTCCCAGATCAGTTTTACAGGTCTGGCTGTCTCTTTTCCTACATGTTGGGTAA GTTCCACACAACCCGACGACGGAAGGAGTTGGGACTCCAGGGTCTAGTCAAAATGTCCAGACCGACAGAGAAAAAGGATGTACAACCCATT K V C W A A F L N P E V P D O F Y R S G C L F S Y M L G K R S S M P P N P T P V M D T Q A D P W G K V P G P G Y G R S NLPKTTAPEVSG. (SCOJD NO:32) (SEGIDNO: F TGTCTTGTGGGTCTTTTCAAATGACATCCCTGAAGGGGACCTGGAGGAAGGTGGTCCGGCTGCACCCTATCGC 704 ACAGAACACCCAGAAAAGTTTACTGTAGGGACTTCCCCTGGACCTCCTTCCACCAGGCCGACCGTGGGATAGCG

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

on page 13 ,-line-	Further deposits are identified on an additional sheet
	runner deposits are identified on an additional sneet
Name of depositary institution American Type Culture Collection	·
Address of depositary institution (including postal code and	(country)
12301 Parklawn Drive Rockville, Maryland 20852 United States of America	
Date of deposit May 16, 1997	Accession Number
Date of deposit May 16, 1997	209053
C. ADDITIONAL INDICATIONS (leave blank if not app	olicable) This information is continued on an additional sheet
D. DESIGNATED STATES FOR WHICH INDICATION	NS ARE MADE (if the indications are not for all designated States)
D. DESIGNATED STATES FOR WHICH INDICATION	NS ARE MADE (if the indications are not for all designated States)
D. DESIGNATED STATES FOR WHICH INDICATION	NS ARE MADE (if the indications are not for all designated States)
D. DESIGNATED STATES FOR WHICH INDICATION	NS ARE MADE (if the indications are not for all designated States)
SEPARATE FURNISHING OF INDICATIONS (leave	e blank if not applicable)
. SEPARATE FURNISHING OF INDICATIONS (leave	e blank if not applicable)
. SEPARATE FURNISHING OF INDICATIONS (leave the indications listed below will be submitted to the Internation	e blank if not applicable)
. SEPARATE FURNISHING OF INDICATIONS (leave the indications listed below will be submitted to the Internation	e blank if not applicable)
. SEPARATE FURNISHING OF INDICATIONS (leave the indications listed below will be submitted to the Internation tumber of Deposit")	e blank if not applicable) onal Bureau later (specify the general nature of the indications e.g., "Accessic
separate furnishing of indications (leave the indications listed below will be submitted to the Internation tumber of Deposit") For receiving Office use only	e blank if not applicable) onal Bureau later (specify the general nature of the indications e.g., "Accessic For International Bureau use only
. SEPARATE FURNISHING OF INDICATIONS (leave the indications listed below will be submitted to the Internation tumber of Deposit")	e blank if not applicable) onal Bureau later (specify the general nature of the indications e.g., "Accessio For International Bureau use only
SEPARATE FURNISHING OF INDICATIONS (leave the indications listed below will be submitted to the International fumber of Deposit") For receiving Office use only	e blank if not applicable) onal Bureau later (specify the general nature of the indications e.g., "Accessio For International Bureau use only

REC'D

2 3 FED 1998

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

	WIPO PCT
A. The indications made below relate to the microorganism refer	ed to in the description
on page <u>13</u> . ime <u>T</u>	able l
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution	
American Type Culture Collection	
Address of depositary institution (including postal code and count	איר
12301 Parklawn Drive Rockville, Maryland 20852	
United States of America	
	•
Date of deposit	Accession Number
May 16, 1997	20 905 3
C. ADDITIONAL INDICATIONS (leave blank if not applicable	This information is continued on an additional sheet
EUROPE	
In respect of those designations in which	a European Patent is sought a sample
of the deposited microorganism will be ma	de available until the publication of
the mention of the grant of the European application has been refused of withdrawn	patent or until the date on which the
the issue of such a sample to an expert n	
sample (Rule 28(4) EPC).	
D. DESIGNATED STATES FOR WHICH INDICATIONS AF	LE MADE (if the indications are not for all designated States)
E. SEPARATE FURNISHING OF INDICATIONS (leave blan.	k if not applicable)
The indications listed below will be submitted to the International B Number of Deposit")	ureau later (specify the general nature of the indications e.g., "Accession
Number of Deposit)	
·	
. 84.	
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	This sheet was received by the International Bureau on:
7 ms sheet was received with the international application	This sheet was received by the international Bureau on.
Authorized offices verte simples	Authorized officer
Particuel Specialist	
1450-501 Operations	1

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorgan	
on page 13 ,lihd,	/ Table 1
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code of 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	and country)
Date of deposit May 16, 1997	Accession Number 209054
C. ADDITIONAL INDICATIONS (leave blank if not	applicable) This information is continued on an additional sheet
D. DESIGNATED STATES FOR WHICH INDICAT	FIONS ARE MADE (if the indications are not for all designated States)
E. SEPARATE FURNISHING OF INDICATIONS (In the indications listed below will be submitted to the internal Number of Deposit")	leave blank if not applicable) mational Bureau later (specify the general nature of the indications e.g., "Accession
	**
This sheet was received with the international applic Authorized officer Authorized Specialist APD-FCT Operations APD-FCT/RO/134 (July 1992)	For International Bureau use only Cation This sheet was received by the International Bureau on: Authorized officer

59/2

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

2 3 FEB 1998 REC'D PCT WiPO

A. The indication	ns made below relate to the mid	roorganism refe	erred to in the description
on page	13	,-line Ta	ble l
B. IDENTIFICA	LTION OF DEPOSIT		Further deposits are identified on an additional sheet
Name of depositar	y institution		
America	a Type Culture Coll	ection	
12301 Pa Rockvil	tary institution (including posta arklawn Drive Le, Maryland 20852 States of America	al code and cour	ntry)
Date of deposit	May 16, 1997		Accession Number 209054
C. ADDITIONA	L INDICATIONS (leave blan	k if not applicab	ble) This information is continued on an additional sheet
the mention application by the issu the sample	of the grant of the has been refused the of such a sample (Rule 28(4) EPC).	m will be he Europea or withdra to an exp	ich a European Patent is sought a sample made available until the publication on patent or until the date on which awn or is deemed to be withdrawn, only pert nominated by the person requesting RE MADE (if the indications are not for all designated States)
E. SEPARATE F	URNISHING OF INDICATI	ONS (leave blas	nk if not applicable)
The indications list Number of Deposit")	ed below will be submitted to the	e International I	Buteau later (specify the general nature of the indications e.g., "Accession
	or receiving Office use only		For International Bureau use only
This sheet was	received with the internationa	l application	This sheet was received by the International Bureau on:
Authorized officer	Paralegal Specialis IAPD-PCT Operati	oue oue	Authorized officer

15

20

25

What is claimed is:

An isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y; or a nucleotide sequence complementary to said isolated polynucleotide.

60

- 2. The polynucleotide of claim 1 wherein said polynucleotide comprises the nucleotide sequence contained in SEQ ID NO:X encoding the receptor polypeptide of SEQ ID NO:Y.
 - 3. The polynucleotide of claim 1 wherein said polynucleotide comprises a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length.
- 4. The polynucleotide of claim 3 which is polynucleotide of SEQ ID NO: X.
 - 5. The polynucleotide of claim 1 which is DNA or RNA.
- 6. A DNA or RNA molecule comprising an expression system, wherein said expression system is capable of producing a receptor polypeptide comprising an amino acid sequence, which has at least 80% identity with the polypeptide of SEQ ID NO:Y when said expression system is present in a compatible host cell.
 - 7. A host cell comprising the expression system of claim 6.
- 8. A process for producing a receptor polypeptide comprising culturing a host of claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.
- A process for producing a cell which produces a receptor polypeptide thereof comprising transforming or transfecting a host cell with the expression system of claim 6 such that the host cell, under appropriate culture conditions, produces a
 receptor polypeptide.
 - 10. A receptor polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO:Y over its entire length.

11. The polypeptide of claim 10 which comprises the amino acid sequence of SEQ ID NO:Y.

61

- 5 12. An antibody immunospecific for the receptor polypeptide of claim 10.
 - 13. A method for the treatment of a subject in need of enhanced activity or expression of receptor polypeptide of claim 10 comprising:
- (a) administering to the subject a therapeutically effective amount of an agonist to said receptor; and/or
 - (b) providing to the subject polynucleotide of claim 1 in a form so as to effect production of said receptor activity in vivo.
- 14. A method for the treatment of a subject having need to inhibit activity or expression of the receptor polypeptide of claim 10 comprising:
 - (a) administering to the subject a therapeutically effective amount of an antagonist to said receptor; and/or
 - (b) administering to the subject a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said receptor; and/or
 - (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said receptor for its ligand.
 - 15. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of the receptor polypeptide of claim 10 in a subject comprising:
 - (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said receptor polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of the receptor polypeptide
 30 expression in a sample derived from said subject.
 - 16. A method for identifying agonists to the receptor polypeptide of claim 10 comprising:
 - (a) contacting cells produced by claim 9 with a candidate compound;
- 35 and

20

25

(b) determining whether the candidate compound effects a signal generated by activation of the receptor polypeptide.

15

- 17. An agonist identified by the method of claim 16.
- 18. The method for identifying antagonists to the receptor polypeptide of claim 10 comprising:

62

- (a) contacting said cell produced by claim 9 with an agonist; and
- (b) determining whether the signal generated by said agonist is diminished in the presence of a candidate compound.
 - 19. An antagonist identified by the method of claim 18.

20. An isolated receptor polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC; and
- (b) a nucleotide sequence complementary to the nucleotide sequence of (a).
- 21. A recombinant host cell produced by a method of Claim 9 or a membrane thereof expressing a receptor polypeptide.

20 30 40	RERVEGOKSNRKDYSLTMOSSVTVOEGM HMACR70.AA GSLOEKPVYELOVOKSVTVOEGL OB-1.aa	VDSQTDSDPVHGYWFRAGNDISWKAPVAT HMACR70.AA WRSWYSSPPLYVYWFRDGEIPYYAEVVAT OB-1.aa	RDRFHLLGDVOKKNCSLSIGDARMEDTGSOB-1.2a	KWNYKYDOLSVHAPCR70.AA	180 200
10	CC	S S S Y P C S F S Y P	NNPAWAVQEETRDRF NNPDRRVKPETQGRF	130 YFFRMEKG-NIKWN YFFRVERGRDVKYS	ıs
		4 %	8 %	121	142

F16.1A

		210 1	220 1	230	240
142	SELTLTPRP	H O E	 QMKRQGAQVT	GTNLTCQMKRQGAQVTTERTVQLNVSYAOB-1.aa	Y P HMACR70.AA
		250	260	270	280
147	PONLTVTV POTITI	FOGEGTASTA FR-NGIALEI	LGNSSSLSVLLONTSYLPVE	VLEGOSLRIVCAVD HACR70.AA VEEGOALRLLCDAP OB-1.aa	VD HMACR70.AA
		290	300	310	320
187	SNPPARLSW SNPPAHLSW	WR SLTI	YPSOPSNPLVLEI NATPISNTGILEL	LELQ-VHLGDEGEFHMACR70 LELRRVRSAEEGGFOB-1.aa	E F HWACK70.AA G F OB-1.aa
		330	340	350 1	360
313	TCRAQNSLGS TCRAQHPLGF	QHVSLNLS LQIFLNLS	VYSLPQLLGP	SCSWEAEGLHCR	HMCR70.AA C R OB-1.aa
		370 1	380	390	400
244 353	CSFRARPAPS	 LCWRLEEK	T T T T T T T T T T T T T T T T T T T	FKVNSSSAGP	HMACR70.AA P W A OB-1.aa

FIG. 18






FIG. 10



F16.2A

SUBSTITUTE SHEET (RULE 26)



F16.2B

MRC-0X44.aa HTEDK48.aa PETA-3.aa A 10 H


245 218 252 HPW∠E25.aa NAG-2.aa TALL∆-1.aa HPW≙E25.aa NAG-2.aa TALLA-1.aa 120 F K D E K K D E K K D E K K D E K K D E K K D E K K D E K K D E K K D E K K D E 0 4 < < > A 17 (2 I G A < \cup >00 V N V L S A A G V A P V L Z Z V JAH SPS 엄마 v m 5 N A L O H SZS ய ய.ம 0 0 F A A A C (J (J <<

38 38

10000

SUBSTITUTE SHEET (RULE 26)


116 113 113	130 140 150 150 160 HFLTLLVVPAIKKDVGSQEDFTQVWNTTMKGLKCC HPWAE25RYAQQDLKKGLHLYGTQGNVGLTNAWSII QTDFRCC NAG-2.a	. aa . aa . aa
151 149 147	170 . 200 .	5. aa aa L. aa
190 180 179	210 220 230 240 240 220 230 230 240 240 240 240 240 240 240 240 240 24	5. aa aa 1. aa
218 213 214	250 270 280 V A A G I G - G L E L A - A M N C V H V S V L Q S T I S P L L P L P L L L HPWAE25. 3 A L V Q I L - G L T F A M T M Y C	5.aa aa 1.aa
253 229 234	290 300 HPWAE25. 3 P H G N C E E A P W Q A A V I G G G D R - I NAG-2.aa 9 0 V V K A D T Y - C A TALLA-1. 4 R F I T A N Q Y E M V	5.aa aa 1.aa



F16.4A

HTPEF86.aa B1.aa	HTPEF86.aa Bl.aa	HTPEF86.aa Bl.aa	HIPEF86.aa B1.aa
240 EFGI IAGI VEN	270	320 PQDQESSP	
230 C L L E A F F Q E L V I	270 I Y P N I Y A A N I E I K E	310 NFPEP	
220 1 S G V L L V F 1 L S V M L I F	CCQSSNVSVI SAEEKKEQTI	300 	
210 	250 CASSHFGCQLV SRPKSNIVLL	290 PSYSSEIQ PKNEEDIEIIPI	አ S ዓ P
182 175 KNSPS	199 A 215 EWKRT	235 PVTSP 251 ETSSQ	248 ANK 291 IENDSS

FIG. 4B



F16.5A




FIG. 5B




FIG. 6A




FIG.6B

	MARACLOAVKYLM		PWLGGCGVI	FAFNLFFWLGGCGVLGVGIWLAATQGS	
	MANACLUAVA 1			1 C V G I W L A A T Q.	6 S NAU-2.aa
44	FATLSSSFPS	LSALSA	LTGAFVMAIC TGAFVMAIC	ANLLIITGAFVMAIGFVGCLGAIKENK HTPBA27. ANLLIITGAFVMAIGFVGCLGAIKENK NAG-2.aa	N K HTPBA27.aa N K NAG-2.aa
	- 8	0	100	110	120
छ छ	CLLLTFFLLLLLV	_	FIATLFFAY1 FIAILFFAY1	F L L E A T I A I L F F A Y T D K I D R Y A Q Q D L K F L L E A T I A I L F F A Y T D K I D R Y A Q Q D L K	LK HTPBA27.aa LK NAG-2.aa
121	KGLHLYGTOGNV KGLHLYGTOGNV	U U	140 SIIOTDFRCC SIIOTDFRCC	LTNAWSIIOTDFRCCGVSNYTDWFEVY HTPRA27.aa	160 V Y HTPBA27.aa V Y NPG-2.aa
161	NATRVPDSCC	17 th 18	180 HAPGTWWKA	SESCGLHAPGTWWKAPCYETVKVWLOE	200 O.E. HTPEA27.aa

FIG. 7A




FIG.7B




FIG.8A




FIG.8B

HHFEK40.aa PETA-3.aa	HHFEK40.aa PETA-3.aa	HHFEK40.aa PETA-3.aa	.60 S HHFEK40.aa N PETA-3.aa
40 A A V L A V G I W A V M A V G I W	NVTGFLGFG	120 HVYYORLSD YAYYOOLNT	160 FKCCGSNS
30 FNFFFWVGG YNCCFWLAGI	SSTFAASAYILIFAGVLVMVTGFLGFG HFFK40.aa SGTYLATAYILVVAGTVMVTGVLGCC PETA-3.aa	110 VELVAGVLA LELIAGILA	150 HASVDRLQQ TSAVDQLQQ0
20 30 40 IYLKYLLFVFNFFFWVGGAVLAVCIW HFFK40.aa VCLKYLFTYNCCFWLAGIW PETA-3.aa		FCLLLVIFLVELVAGVLAHVYYORLSD FILLLIFLLEIIAGILAYAYOQLNT	140 150 160 160 17 G Q P E H A D - H A S V D R L Q Q D F K C C G S N S 18 H Q P G H E A V T S A V D Q L Q Q E F H C C G S N N
10 MAHYKTEQDDWLI MGEFNEKKTTCGT	TLVEKSGYLSVLA TLALKSDYLSLLA	90 AILWERKGCLSTY ATEKERRNLLRLY	130 ELKQHLNRTLAEN ELKENLKDTMTKR
	44	81 81 A	121

FIG.9A




FIG. 9F

19/27

₩ ₩	10 MCTGKCARCVGLSLITLC MCTGKCARCVGLSLITLC	20 LVCIVANALI FVCIVANALI	TLCLVCIVANALLLVPNGETSWTNT	40 T HGBGV89.aa T L6H.aa
4 4	SO NHLSLQVWLMGGFIGGGI NHLSLQVWLMGGFIGGGI	60 MVLCPGIAAV	IGGGLMVLCPGIAAVRAGGKGCCGAGC IGGGLMVLCPGIAAVRAGGKGCCGAGC	80 C HGBGV89.aa C L6H.aa
ਲ ਲ	CGNRCRMLRSVFSSAFGVCGNRCRMLRSVFSSAFGV	TOO FGVLGAIYCLSVS FGVLGAIYCLSVS	SAFGVLGAIYCLSVSGAGLRNGPRCLMSAFGVLGAIYCLSVSGAGLRNGPRCLM	120 M HGBGV89.aa M L6H.aa
121	130 NGEWGYHFEDTAG NGEWGYHFEDTAG	140 AYLLNRTLWDRCEAI AYLLNRTLWDRCEAI	150 160 PPRVVPWNVTLFS PPRVVPWNVTLFS	160
161	170 LLVAASCLEIVLCGI LLVAASCLEIVLCGI	180 190 0 L V N A T I G V F C G D C R K K Q D T P Q L V N A T I G V F C G D C R K K Q D T P	190 DCRKKODTPH DCRKKODTPH	HGBGV89.aa L6H.aa

F16.10




FIG. 11A




FIG. 11B

HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8.aa
40 40 40 40 40 40 40 40 40 40 40 40 40 4	8 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	120 H r IVIMVL h	160 1 160 1 H S I G
DCSHALPQGPFFTSIPPNGPFVGTPDO	70 KPRADVAFH	110 FOKEKKSFE	150 DTLGIYGKVN
20 V V LONIIYNPVI	VSLRDQAA INLRC-GG VDLQN-GSSM	100 	140 YGHRIGPEKI
10 PRLE- SFSTQTPYPNLA-	VRGLVLQEPKHFT ISGVVLSDAKRFQ IRGHVPSDADRFQ	90 1 	130
222 222	33 33 34 34 34 37 37	53 72 R A A B A	46 112 K A

FIG. 12A

SUBSTITUTE SHEET (RULE 26)

HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rCalectin-5.aa hCalectin-8.aa
200 	240 	ZBO Y P Q R F F E V I S R G Q R F S V	ELRISGSV TEVAGDI TEINGDI	
190 190 190 190 190 190 190 190	230 NVDLLAGK	270 LISAPFLF SLPGSMPF RNITSFPF	310 Q Q A L E Q L R L M N L P D I N F K E L S S I D	2B
180 180 ELTEISREN	220 	260 	300 QGLGATSMN QHICEYSHR VHSLEYKHR	FIG.12B
170 	210 	250 FADRTLAWIS FDENAVVRNTQI LNIKAFVRNSFI	POEGGLKLAUNG CEGHCFKVAVDG CDVREFKVAVNG	330 CVHS. -THVET -LEVRSW
46 53 152 F S F	46 53	52 R A S 59 N P R 232 N P R	87 LLL 99 WIL 271 IIY	127 Q 11 Y 139 Q 11 - 311 H 11 -




FIG. 15

HMMGS46.aa B-cell Recpt Associated Pr.aa	1 60 IGGVQQ HMMCS46.aa IGGVQQ B-cell Recpt Associated Pr.aa	PRKI B-cell Recpt Associated Pr.aa	120 L P S M HMWSS46.aa L P S M B-cell Recpt Associated Pr.aa	 HMMGS46.aa B-cell Recpt Associated Pr.aa
MAQNLKDLAGRLPAGPRGMGTALKLLLGAG HMMSS46.aa	AVAYGVRESVFTVEGGHRAIFFNRAVAYGVRESVFTVEGGHRAIFFNR	70 80 DTILAEGLHFRIPWFQYPIIYDIRAR DTILAEGLHFRIPWFQYPIIYDIRAR	SSPTGSKDLQMVNISLRVLSRPNAQE SSPTGSKDLQMVNISLRVLSRPNAQE	121 YORLGLDYEERVLPSIVNEVLKSVVAKFNA HMGS46.aa 121 YORLGLDYEERVLPSIVNEVLKSVVAKFNA B-cell Rec
44	ਜ਼ ਜ਼	ਰ ਰ	ਲ ਲ	22

FIG. 14A

FIG. 14B

HNFGWD6.aa EGFR related-protein.aa	HNFGAD6.aa EGFR related-protein.aa	HNFGWD6.aa EGFR related-protein.aa	HNFGWD6.aa EGFR related-protein.aa	sd-protein.aa	
				0 FINEGAD6.aa EGFR related-protein	SCANORAL STATES
TREYCEFME SREYCDFMR	EERTLGMAA	90 6 A E H H M P I P	DFYRSGCLF OFYRL	150 A D P W G K V P G W	
20 TKGSCEIT TKGRCEIT	50 50 R G R P G V V H H	80 8 ASDSGCFW 	FLNPEVPD FLNPEVPD	140 P T P V M D T Q	(((((
IKGRPCCIG	40 EEATLCSQV EEATLCSQV	70 1 GSRTPSHVG 	100 LDKVCWAAA MDDVCGLLP	130 G K R S S M P P N	160
M D C E	GYFH	Ø 1 ↔	R C - C - C - C - C - C - C - C -	121 S Y M L 68	

International Application No PCT/#S 98/00959

			101/05 5	0/00333
A. CLASS IPC 6	FICATION OF SUBJECT MATTER C12N15/12 C12N15/85 C12N5/ C12Q1/68 A61K38/17 A61K48		705 C07	K16/28
According t	to International Patent Classification (IPC) or to both national classification	ication and IPC		
B. FIELDS	SEARCHED			
Minimum d IPC 6	ocumentation searched (classification system tollowed by classification C12N C07K C12Q A61K	ation symbols)		
Documenta	tion searched other than minimum documentation to the extent that	such documents are include	ed in the fields se	earched
Electronio d	lata base consulted during the international search (name of data b	ase and, where practical, se	earch terms used)
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT			
Category *	Citation of document, with indication, where appropriate, of the re	elevant passages		Relevant to claim No.
X	DATABASE GENBANK Accession No. T48852, 8 February "EST70319 from Stratagene place XP002066898 see the whole document			1-12,20
X	DATABASE GENBANK Accession No. G14442, 4 January "STS SHGC-10055" XP000206689 see the whole document	1996		1-3,20
		-/		
X Furth	er documents are listed in the continuation of box C.	Patent family men	nbers are listed in	n annex.
"A" document consider the consider of filing da "L" document which is citation "O" document other m "P" document later that the consideration of the administration of the admin	t which may throw doubts on priority claim(s) or cited to establish the publication date of another or other special reason (as specified) at referring to an oral disclosure, use, exhibition or eans at published prior to the international filing date but.	To later document publishing or priority date and no cited to understand the invention. "X" document of particular cannot be considered involve an inventive stand to be considered document for combined ments, such combination the art. "&" document member of the invention of t	tin conflict with the principle or the relevance; the clinovel or cannot the pwhen the doc relevance; the clito involve an involve a	the application but cory underlying the aimed invention be considered to summent is taken alone aimed invention entive step when the cother such docuse to a person skilled amily
	June 1998	Authorized officer	6. 09. 1998	
	European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo ni, Fax: (+31-70) 340-3016	Galli, I		

Form PCT/ISA/210 (second sheet) (July 1992)

Internatic—al Application No
PCT/US 98/00959

0.00======	etion) DOCUMENTS CONSIDERED TO BE RELEVANT	101/03 30/00333
C.(Continue	- A of the minute and a second	Relevant to claim No.
Category	Olabert of Sections with the section of the section	
A .	ZANNETTINO A.C.W. ET AL.: "A powerful technique for isolating genes encoding surface antigens using retroviral expression cloning" J. IMMUNOL., vol. 156, no. 2, 15 January 1996, pages 611-620, XP002066893 see abstract	1-21
A	ADAMS M D ET AL: "COMPEMENTARY DNA SEQUENCING: EXPRESSED SEQUENCE TAGS AND HUMAN GENOME PROJECT" SCIENCE, vol. 252, no. 5013, 21 June 1991, pages 1651-1656, XP000645049 cited in the application see abstract	1-21
A	SIMMONS D. & SEED B.: "Isolation of a cDNA encoding CD33, a differentiation antigen of myeloid progenitor cells" J. IMMUNOLOGY, vol. 141, no. 8, 15 October 1988, pages 2797-2800, XP002066894 see abstract see figure 4	1-21
A .	CROCKER P.R. ET AL.: "Sialoadhesin, a macrophage sialic acid binding receptor for haemopoietic cells with 17 immunoglobulin-like domains" EMBO J., vol. 13, no. 19, 3 October 1994, pages 4490-4503, XPO02066895 see abstract see figure 2	1-21
A	KELM S. ET AL.: "The Sialoadhesins: a family of sialic acid-dependent cellular recognition molecules within the immunoglobulin superfamily" GLYCOCONJ. J., vol. 13, no. 6, December 1996, pages 913-926, XP002066896 see the whole document	1-21
A	DATABASE GENBANK Accession No. U71382, 12 November 1996 PATEL N. ET AL.: "OB binding protein - 1" XP002066901 cited in the application see the whole document -/	1-21
	-/	

International Application No PCT/US 98/00959

10-	PCT/US 98/00959	
.(Continu:	ation) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication; where appropriate, of the relevant passages	Relevant to claim No.
P,X	DATABASE GENBANK Accession No. AA344713, 21 April 1997 "EST50650, H. sapiens cDNA similar to CD33 antigen" XP002066900 see the whole document	1-12,20
P,A	TAKEI Y. ET AL.: "Molecular cloning of a novel gene similar to myeloid antigen CD33 and its specific expression in placenta" CYTOGENET. CELL GENET., vol. 78, 1997, pages 295-300, XP002066897 see abstract see figure 1	1-21
·		
	-9	

nter onal application No.

PCT/US 98/00959

Box (Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This Inte	ernational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. X 2.	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely: Remark: Although claims 13 and 14 are directed to a method of treatment of the human/animal body, and claim 15 (as far as in vivo methods are envisaged) is directed to a diagnostic method, the search has been carried out and based on the alleged effects of the compound/composition. Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3.	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II	Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)
	rnational Searching Authority found multiple inventions in this international application, as follows:
	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
,	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: see further information sheet, subject 1.
Remark c	The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: (1-21) - partial

An isolated polynucleotide encoding the HMACR70 cell surface receptor (Seq. IDs 1,18), polynucleotides complementary thereto, fragments thereof, homologs thereof.

An expression system comprising said polynucleotides, a process to produce a cell harbouring said expression system. Said cell or membrane thereof expressing said receptor.

A process for the production of the HMACR70 polypeptide comprising culturing said cell.

Said HMACR70 polypeptide or homologs. An antibody directed against said polypeptide.

Therapeutic and diagnostic applications.

A method to identify agonists or antagonists to said receptors. Said agonists and antagonists.

2. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 2,3,19

3. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 4,20

4. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 5,21

5. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 6,22

6. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 7,23

7. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 8,24

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

- 8. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 9,10,25
- 9. Claims: (1-21) partial

 Idem as subject matter 1 but limited to Seq. IDs 11,26
- 10. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 12,27
- 11. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 13,28
- 12. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 14,29
- 13. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 15,30
- 14. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 16,31
- 15. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 17,21

